

UPPER BOUNDS ON THE SLOPE OF CERTAIN FIBERED SURFACES

MAKOTO ENOKIZONO

ABSTRACT. We establish the slope equality and give an upper bound of the slope for finite cyclic covering fibrations of an elliptic surface including bielliptic fibrations of genus greater than 5. We also give an upper bound of the slope for triple cyclic covering fibrations of a ruled surface and hyperelliptic fibrations, which provides a new proof of Xiao's upper bound.

INTRODUCTION

We shall work over the complex number field $\mathbb{C}$. Let S be a smooth projective surface, B a smooth projective curve and $f: S \rightarrow B$ a relatively minimal fibration of curves of genus $g \geq 2$. Let K_f denote the relative canonical bundle $K_S - f^*K_B$. Consider the following relative invariants:

$$\begin{aligned}\chi_f &:= \chi(\mathcal{O}_S) - \chi(\mathcal{O}_B)\chi(\mathcal{O}_F), \\ e_f &:= e(S) - e(B)e(F),\end{aligned}$$

where F denotes a general fiber of f and $e(M)$ the topological Euler characteristic of M . Then the following are well known:

- (Noether) $12\chi_f = K_f^2 + e_f$.
- (Arakelov) K_f is nef.
- (Ueno) $\chi_f \geq 0$, and $\chi_f = 0$ if and only if f is locally trivial (i.e., a holomorphic fiber bundle).
- (Segre) $e_f \geq 0$, and $e_f = 0$ if and only if f is smooth.

Assume that f is not locally trivial. We put

$$\lambda_f = \frac{K_f^2}{\chi_f}$$

and call it the *slope* of f . Then one sees $0 < \lambda_f \leq 12$ from the above results. In 1987, Xiao showed in [17] the inequality

$$\lambda_f \geq 4 - \frac{4}{g}.$$

This inequality is sharp and it is well known that the slope attains the lower bound only if the fibration f is hyperelliptic, i.e., a general fiber F is a hyperelliptic curve ([13] and [17]). As to the upper bound, Kodaira [11] constructed examples of fibrations with slope 12, which are nowadays called Kodaira fibrations. Thus the inequality $\lambda_f \leq 12$ is sharp among all

2010 *Mathematics Subject Classification.* 14D06.

Keywords: fibered surface, slope inequality, cyclic covering.

fibered surfaces. On the other hand, Matsusaka [16] obtained an upper bound smaller than 12 for hyperelliptic case and Xiao [19] improved this bound. In [5], upper bounds for genus 3 fibrations are studied from another point of view.

We studied in [7] primitive cyclic covering fibrations of type (g, h, n) . Roughly speaking, it is a fibered surface of genus g obtained as the relatively minimal model of an n sheeted cyclic branched covering of another fibered surface of genus h . Note that hyperelliptic fibrations are nothing more than such fibrations of type $(g, 0, 2)$ and that bielliptic fibrations of genus $g \geq 6$ are those of type $(g, 1, 2)$ (cf. [3] and [6]). Here, a fibration is called *bielliptic* if a general fiber is a bielliptic curve, i.e., a non-singular projective curve obtained as a double covering of an elliptic curve. In [7], we established the lower bound of the slope for such fibrations of type (g, h, n) extending former results for $n = 2$ in [2] and [6]. Furthermore, when $h = 0$ and $n \geq 4$, we obtained even the upper bound (expressed as a function in g and n) which is strictly smaller than 12. Recall that known examples of Kodaira fibrations, including Kodaira's original ones [11], are presented as primitive cyclic covering fibrations with $h \geq 2$ and, in fact, there exist such fibrations for any $h \geq 2$ (see, [4] and [10]). Hence, as far as the upper bound of the slope strictly smaller than 12 concerns, the remaining cases to be examined are $h = 0, n = 3$ and $h = 1, n \geq 2$.

The purpose of the present article is to give an affirmative answer to the above mentioned upper bound problem by introducing numerical invariants attached to the singularities of the branch locus of the cyclic covering, and improving a coarser estimates in [7]. When $h = 0$, a vertical component of the branch locus on a relatively minimal model is always a non-singular rational curve and this fact makes it much easier to handle singularities on the branch locus. On the other hand, when $h \geq 0$, we must pay attention to all subcurves of fibers and their singularities in a fibration of genus h , which seems quite terrible. Fortunately enough, when $h = 1$, we have Kodaira's classification of singular fibers [12] from which we know that major components are rational curves and singularities are mild. This gives us a hope to extend results for $h = 0$ to fibrations of type $(g, 1, n)$. In fact, we can show the following:

Theorem 0.1. *Assume that $g \geq (2n - 1)(3n - 1)/(n + 1)$. Then, there exists a function $\text{Ind}: \mathcal{A}_{g,1,n} \rightarrow \mathbb{Q}_{\geq 0}$ from the set $\mathcal{A}_{g,1,n}$ of all fiber germs of primitive cyclic covering fibrations of type $(g, 1, n)$ such that $\text{Ind}(F_p) = 0$ for a general $p \in B$ and*

$$K_f^2 = \frac{12(n-1)}{2n-1} \chi_f + \sum_{p \in B} \text{Ind}(F_p)$$

for any primitive cyclic covering fibration $f: S \rightarrow B$ of type $(g, 1, n)$.

Theorem 0.2. *Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(g, 1, n)$. Then*

$$\lambda_f \leq 12 - \begin{cases} \frac{6n^2}{(n+1)(g-1)}, & \text{if } n \geq 4, \text{ or } n = 3 \text{ and } g = 4 \\ \frac{24}{4g-17}, & \text{if } n = 3 \text{ and } g > 4, \\ \frac{2}{g-2}, & \text{if } n = 2 \text{ and } g \geq 3. \end{cases}$$

In particular, we have the slope equality and the upper bound of the slope for bielliptic fibrations of genus $g \geq 6$. We remark that the upper bounds in Theorem 0.2 are “fiberwise” sharp as we shall see in Example 3.13. We do not know, however, whether there exist primitive cyclic covering fibrations of type $(g, 1, n)$ whose slopes attain the bounds.

It remains to investigate the case that $h = 0$ and $n \leq 3$. The above method also applies to this case and we obtain the following:

Theorem 0.3. *Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(g, 0, n)$*

- (1) *If $g = 4$ and $n = 3$, then $\lambda_f \leq 129/17$ with the equality sign holding if and only if any singular fiber of f is a triple fiber.*
- (2) *Assume that $n = 3$. If either $g > 4$, or $g = 4$ and f has no triple fibers, then*

$$\lambda_f \leq 12 - \frac{72(g+1)}{4g^2 + g + 13 - 36\delta},$$

where $\delta = 0$ if $g + 2 \in 6\mathbb{Z}$ and $\delta = 1$ otherwise.

- (3) *If $n = 2$, then*

$$\lambda_f \leq 12 - \frac{4(2g+1)}{g^2 - 1 + \delta} = 12 - \frac{4(2g+1)}{g^2 + \frac{(-1)^g - 1}{2}},$$

where $\delta = 0$ if g is odd and $\delta = 1$ if g is even, i.e., $2\delta = 1 + (-1)^g$.

We obtained the complete classification of singular fibers of primitive cyclic covering fibrations of type $(4, 0, 3)$ in [8] and computed some local invariants for each singular fiber. One may use it to reprove Theorem 0.3 (1). Note also that Theorem 0.3 (3) provides a new proof of Xiao’s upper bound for hyperelliptic fibrations in [19] referred above. It is shown in [15] that the inequality is optimal for given g .

The organization of the paper is as follows. In §1, we recall basic results from [7] and [8] on primitive cyclic covering fibrations and introduce some notation for the later use. In §2, we observe the local concentration of relative invariants of primitive cyclic covering fibrations of type $(g, 1, n)$ on a finite number of fiber germs and show Theorem 0.1. §3 will be devoted to the proof of Theorem 0.2. In the course of the study, we freely use Kodaira’s table of singular fibers of elliptic surfaces. Finally in §4, we show Theorem 0.3.

Acknowledgment. The author expresses his sincere gratitude to Professor Kazuhiro Konno for many suggestions and warm encouragement.

1. PRELIMINARIES

In this section, we recall and state basic results for primitive cyclic covering fibrations in [7].

Definition 1.1. A relatively minimal fibration $f: S \rightarrow B$ of genus $g \geq 2$ is called a primitive cyclic covering fibration of type (g, h, n) , if there exist a (not necessarily relatively minimal)

fibration $\tilde{\varphi}: \tilde{W} \rightarrow B$ of genus $h \geq 0$, and a classical n -cyclic covering

$$\tilde{\theta}: \tilde{S} = \operatorname{Spec}_{\tilde{W}} \left(\bigoplus_{j=0}^{n-1} \mathcal{O}_{\tilde{W}}(-j\tilde{\mathfrak{d}}) \right) \rightarrow \tilde{W}$$

branched over a smooth curve $\tilde{R} \in |n\tilde{\mathfrak{d}}|$ for some $n \geq 2$ and $\tilde{\mathfrak{d}} \in \operatorname{Pic}(\tilde{W})$ such that f is the relatively minimal model of $\tilde{f} := \tilde{\varphi} \circ \tilde{\theta}$.

Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type (g, h, n) . We freely use the notation in Definition 1.1. Let $\tilde{F}$ and $\tilde{\Gamma}$ be general fibers of $\tilde{f}$ and $\tilde{\varphi}$, respectively. Then the restriction map $\tilde{\theta}|_{\tilde{F}}: \tilde{F} \rightarrow \tilde{\Gamma}$ is a classical n -cyclic covering branched over $\tilde{R} \cap \tilde{\Gamma}$. Since the genera of $\tilde{F}$ and $\tilde{\Gamma}$ are g and h , respectively, the Hurwitz formula gives us

$$(1.1) \quad r := \tilde{R}\tilde{\Gamma} = \frac{2(g-1-n(h-1))}{n-1}.$$

Note that r is a multiple of n . Let $\tilde{\sigma}$ be a generator of $\operatorname{Aut}(\tilde{S}/\tilde{W}) \simeq \mathbb{Z}/n\mathbb{Z}$ and $\rho: \tilde{S} \rightarrow S$ the natural birational morphism. By assumption, $\operatorname{Fix}(\tilde{\sigma})$ is a disjoint union of smooth curves and $\tilde{\theta}(\operatorname{Fix}(\tilde{\sigma})) = \tilde{R}$. Let $\varphi: W \rightarrow B$ be a relatively minimal model of $\tilde{\varphi}$ and $\tilde{\psi}: \tilde{W} \rightarrow W$ the natural birational morphism. Since $\tilde{\psi}$ is a succession of blow-ups, we can write $\tilde{\psi} = \psi_1 \circ \cdots \circ \psi_N$, where $\psi_i: W_i \rightarrow W_{i-1}$ denotes the blow-up at $x_i \in W_{i-1}$ ($i = 1, \dots, N$) with $W_0 = W$ and $W_N = \tilde{W}$. We define reduced curves R_i on W_i inductively as $R_{i-1} = (\psi_i)_* R_i$ starting from $R_N = \tilde{R}$ down to $R_0 =: R$. We also put $E_i = \psi_i^{-1}(x_i)$ and $m_i = \operatorname{mult}_{x_i}(R_{i-1})$ for $i = 1, 2, \dots, N$.

Lemma 1.2. *With the above notation, the following hold for any $i = 1, \dots, N$.*

- (1) *Either $m_i \in n\mathbb{Z}$ or $m_i \in n\mathbb{Z} + 1$. Moreover, $m_i \in n\mathbb{Z}$ holds if and only if E_i is not contained in R_i .*
- (2) *$R_i = \psi_i^* R_{i-1} - n \left\lceil \frac{m_i}{n} \right\rceil E_i$, where $[t]$ is the greatest integer not exceeding t .*
- (3) *There exists $\mathfrak{d}_i \in \operatorname{Pic}(W_i)$ such that $\mathfrak{d}_i = \psi_i^* \mathfrak{d}_{i-1}$ and $R_i \sim n\mathfrak{d}_i$, $\mathfrak{d}_N = \tilde{\mathfrak{d}}$.*

Let E be a (-1) -curve on a fiber of $\tilde{f}$. If E is not contained in $\operatorname{Fix}(\tilde{\sigma})$, then $L := \tilde{\theta}(E)$ is a (-1) -curve and $\tilde{\theta}^* L$ is the sum of n disjoint (-1) -curves containing E . Contracting them and L , we may assume that any (-1) -curve on a fiber of $\tilde{f}$ is contained in $\operatorname{Fix}(\tilde{\sigma})$. Then $\tilde{\sigma}$ induces an automorphism σ of S over B and ρ is the blow-up of all isolated fixed points of σ (cf. [7]). One sees easily that there is a one-to-one correspondence between $(-k)$ -curves contained in $\operatorname{Fix}(\tilde{\sigma})$ and $(-kn)$ -curves contained in $\tilde{R}$ via $\tilde{\theta}$. Hence, the number of blow-ups in ρ is that of vertical $(-n)$ -curves contained in $\tilde{R}$.

From Lemma 1.2, we have

$$(1.2) \quad K_{\tilde{\varphi}} = \tilde{\psi}^* K_{\varphi} + \sum_{i=1}^N \mathbf{E}_i,$$

$$(1.3) \quad \tilde{\mathfrak{d}} = \tilde{\psi}^* \mathfrak{d} - \sum_{i=1}^N \left[\frac{m_i}{n} \right] \mathbf{E}_i,$$

where $\mathbf{E}_i$ denotes the total transform of E_i . Since

$$K_{\tilde{S}} = \tilde{\theta}^* (K_{\tilde{W}} + (n-1)\tilde{\mathfrak{d}})$$

and

$$\chi(\mathcal{O}_{\tilde{S}}) = n\chi(\mathcal{O}_{\tilde{W}}) + \frac{1}{2} \sum_{j=1}^{n-1} j\tilde{\mathfrak{d}}(j\tilde{\mathfrak{d}} + K_{\tilde{W}}),$$

we get

$$(1.4) \quad K_{\tilde{f}}^2 = n(K_{\tilde{\varphi}}^2 + 2(n-1)K_{\tilde{\varphi}}\tilde{\mathfrak{d}} + (n-1)^2\tilde{\mathfrak{d}}^2),$$

$$(1.5) \quad \chi_{\tilde{f}} = n\chi_{\tilde{\varphi}} + \frac{1}{2} \sum_{j=1}^{n-1} j\tilde{\mathfrak{d}}(j\tilde{\mathfrak{d}} + K_{\tilde{\varphi}}).$$

Definition 1.3 (Singularity index α). Let k be a positive integer. For $p \in B$, we consider all the singular points (including infinitely near ones) of R on the fiber Γ_p of $\varphi: W \rightarrow B$ over p . We let $\alpha_k(F_p)$ be the number of singular points of multiplicity either kn or $kn+1$ among them, and call it the k -th singularity index of F_p , the fiber of $f: S \rightarrow B$ over p . Clearly, we have $\alpha_k(F_p) = 0$ except for a finite number of $p \in B$. We put $\alpha_k = \sum_{p \in B} \alpha_k(F_p)$ and call it the k -th singularity index of f .

Let D_1 be the sum of all $\tilde{\varphi}$ -vertical $(-n)$ -curves contained in $\tilde{R}$ and put $\tilde{R}_0 = \tilde{R} - D_1$. We denote by $\alpha_0(F_p)$ the ramification index of $\tilde{\varphi}|_{\tilde{R}_0}: \tilde{R}_0 \rightarrow B$ over p , that is, the ramification index of $\tilde{\varphi}|_{(\tilde{R}_0)_h}: (\tilde{R}_0)_h \rightarrow B$ over p minus the sum of the topological Euler number of irreducible components of $(\tilde{R}_0)_v$ over p . Then $\alpha_0(F_p) = 0$ except for a finite number of $p \in B$, and we have

$$\sum_{p \in B} \alpha_0(F_p) = (K_{\tilde{\varphi}} + \tilde{R}_0)\tilde{R}_0$$

by definition. We put $\alpha_0 = \sum_{p \in B} \alpha_0(F_p)$ and call it the 0-th singularity index of f .

Let $\varepsilon(F_p)$ be the number of (-1) -curves contained in $\tilde{F}_p$, and put $\varepsilon = \sum_{p \in B} \varepsilon(F_p)$. This is no more than the number of blowing-ups appearing in $\rho: \tilde{S} \rightarrow S$.

From (1.2) and (1.3), we have

$$\begin{aligned}
(K_{\tilde{\varphi}} + \tilde{R})\tilde{R} &= \left(\tilde{\psi}^*(K_{\varphi} + R) + \sum_{i=1}^N \left(1 - n \left\lfloor \frac{m_i}{n} \right\rfloor \right) \mathbf{E}_i \right) \left(\tilde{\psi}^*R - n \left\lfloor \frac{m_i}{n} \right\rfloor \mathbf{E}_i \right) \\
&= (K_{\varphi} + R)R - \sum_{i=1}^N n \left\lfloor \frac{m_i}{n} \right\rfloor \left(n \left\lfloor \frac{m_i}{n} \right\rfloor - 1 \right) \\
(1.6) \quad &= (K_{\varphi} + R)R - n \sum_{k \geq 1} k(nk - 1)\alpha_k.
\end{aligned}$$

On the other hand, we have

$$(1.7) \quad (K_{\tilde{\varphi}} + \tilde{R})\tilde{R} = (K_{\tilde{\varphi}} + \tilde{R}_0)\tilde{R}_0 + D_1(K_{\tilde{\varphi}} + D_1) = \alpha_0 - 2\varepsilon.$$

Hence,

$$(1.8) \quad (K_{\varphi} + R)R = n \sum_{k \geq 1} k(nk - 1)\alpha_k + \alpha_0 - 2\varepsilon.$$

by (1.6) and (1.7). Since $K_f^2 = K_{\tilde{f}}^2 + \varepsilon$, $\chi_{\tilde{f}} = \chi_f$, (1.2), (1.3), (1.4) and (1.5), we get

$$(1.9) \quad K_f^2 = nK_{\varphi}^2 + 2(n-1)K_{\varphi}R + \frac{(n-1)^2}{n}R^2 - \sum_{k \geq 1} ((n-1)k - 1)^2\alpha_k + \varepsilon$$

and

$$(1.10) \quad \chi_f = n\chi_{\varphi} + \frac{(n-1)(2n-1)}{12n}R^2 + \frac{n-1}{4}K_{\varphi}R - \frac{n(n-1)}{12} \sum_{k \geq 1} ((2n-1)k^2 - 3k)\alpha_k.$$

From (1.8), (1.9), (1.10) and Noether's formula, we have

$$(1.11) \quad e_f = ne_{\varphi} + n \sum_{k \geq 1} \alpha_k + (n-1)\alpha_0 - (2n-1)\varepsilon.$$

We define some notation for the later use. For a vertical divisor T and $p \in B$, we denote by $T(p)$ the greatest subdivisor of T consisting of components of the fiber over p . Then $T = \sum_{p \in B} T(p)$. We consider a family $\{L^i\}_i$ of vertical irreducible curves in $\tilde{R}$ over p satisfying:

(i) L^1 is the proper transform of an irreducible curve Γ^1 contained in the fiber Γ_p or a (-1) -curve E^1 appearing in $\tilde{\psi}$.

(ii) For $i \geq 2$, L^i is the proper transform of an irreducible curve Γ^i contained in the fiber Γ_p intersecting Γ^k for some $k < i$ or an exceptional (-1) -curve E^i that contracts to a point x^i on C^k (or on its proper transform) for some $k < i$, where we define C^j to be E^j or Γ^j according to whether L^j is the proper transform of which curve.

(iii) $\{L^i\}_i$ is the largest among those satisfying (i) and (ii).

The set of all vertical irreducible curves in $\tilde{R}$ over p is decomposed into the disjoint union of such families uniquely. We denote it as

$$\tilde{R}_v(p) = D^1(p) + \cdots + D^{\eta_p}(p), \quad D^t(p) = \sum_{k \geq 1} L^{t,k}$$

where η_p denotes the number of the decomposition and $\{L^{t,k}\}_k$ satisfies (i), (ii), (iii). Let $C^{t,k}$ be the exceptional curve or the component of the fiber Γ_p the proper transform of which is $L^{t,k}$. Let $D^t(p)$ be the sum of all irreducible components of $D^t(p)$ which are the proper transforms of curves contained in Γ_p and $D''^t(p) = D^t(p) - D^t(p)$. Let η'_p be the cardinality of the set $\{t = 1, \dots, \eta_p \mid D^t(p) \neq 0\}$ and $\eta''_p = \eta_p - \eta'_p$.

Definition 1.4 (Index j). Let $j_{b,a}(F_p)$ (resp. $j_{b,\bullet}^t(F_p)$, $j_{b,a}^t(F_p)$, $j_{b,a}''^t(F_p)$) be the number of irreducible curves of genus b with self-intersection number $-an$ contained in $\tilde{R}_v(p)$ (resp. $D^t(p)$, $D^t(p)$, $D''^t(p)$). Put

$$j_{\bullet,a}^t(F_p) = \sum_{b \geq 0} j_{b,a}^t(F_p), \quad j_{b,\bullet}^t(F_p) = \sum_{a \geq 0} j_{b,a}^t(F_p), \quad j_{b,a}(F_p) = \sum_{t \geq 1} j_{b,a}^t(F_p).$$

Similarly, we define $j^t(F_p) = j_{\bullet,\bullet}^t(F_p)$, $j_{\bullet,a}^t(F_p)$, $j_{b,\bullet}''^t(F_p)$, etc. Clearly, we have $j_{b,\bullet}''^t(F_p) = 0$ for any $b \geq 1$ by the definition of $D''^t(p)$.

Rearranging the index if necessary, we may assume that $D^t(p) = \sum_{k=1}^{j^t(F_p)} L^{t,k}$, $D''^t(p) = \sum_{k=j^t(F_p)+1}^{j^t(F_p)} L^{t,k}$. Put $L^{t,k} = L^{t,k}$, $L''^{t,k} = L^{t,j^t(F_p)+k}$, $C^{t,k} = C^{t,k}$, $C''^{t,k} = C^{t,j^t(F_p)+k}$.

Let $\alpha_0^+(F_p)$ be the ramification index of $\tilde{\varphi} : \tilde{R}_h \rightarrow B$ over p and put $\alpha_0^-(F_p) = \alpha_0(F_p) - \alpha_0^+(F_p)$. It is clear that $\varepsilon(F_p) = j_{0,1}(F_p)$ and $\alpha_0^-(F_p) = \sum_{b \geq 0} (2b-2)j_{b,\bullet}(F_p) + 2\varepsilon(F_p)$.

Let $\bar{\eta}_p$ be the number of $t = 1, \dots, \eta_p$ such that $j^t(F_p) = j_{0,1}''^t(F_p)$ and $\hat{\eta}_p = \eta''_p - \bar{\eta}_p$.

Definition 1.5 (Vertical type singularity). Let x be a singular point of R . For $t = 1, \dots, \eta_p$ and $u \geq 1$, x is a (t, u) -vertical type singularity or simply a u -vertical type singularity if the number of $C^{t,k}$'s whose proper transforms pass through x is u . If x is a (t, u) -vertical type singularity and the multiplicity of it belongs to $n\mathbb{Z}$ (resp. $n\mathbb{Z} + 1$), we call it a (t, u) -vertical $n\mathbb{Z}$ type singularity (resp. (t, u) -vertical $n\mathbb{Z} + 1$ type singularity).

Let $\iota^{t,(u)}(F_p)$, $\kappa^{t,(u)}(F_p)$ respectively be the number of (t, u) -vertical $n\mathbb{Z}$, $n\mathbb{Z} + 1$ type singularities over p and put

$$\iota^t(F_p) = \sum_{u \geq 1} (u-1)\iota^{t,(u)}(F_p), \quad \kappa^t(F_p) = \sum_{u \geq 1} (u-1)\kappa^{t,(u)}(F_p),$$

$\iota(F_p) = \sum_{t=1}^{\eta_p} \iota^t(F_p)$ and $\kappa(F_p) = \sum_{t=1}^{\eta_p} \kappa^t(F_p)$. Let $\iota_k^{t,(u)}(F_p)$, $\kappa_k^{t,(u)}(F_p)$ respectively be the number of (t, u) -vertical type singularities with multiplicity kn , $kn+1$ and we define $\iota_k^t(F_p)$, $\kappa_k^t(F_p)$, $\iota_k(F_p)$ and $\kappa_k(F_p)$ similarly.

Definition 1.6 (Indices α' , α''). We say that a singular point x of R is involved in $D^t(p)$ if there exists $C^{t,k}$ such that it or its proper transform passes through x or it contracts to x . A singular point x of R is involved in $\tilde{R}_v(p)$ if it is involved in $D^t(p)$ for some t . Let $\alpha'_k(F_p)$ (resp. $\alpha''_k(F_p)$) denotes the number of singularities with multiplicity kn or $kn+1$ over p not

involved in $\tilde{R}_v(p)$ (resp. involved in $\tilde{R}_v(p)$). Clearly, we have $\alpha_k(F_p) = \alpha'_k(F_p) + \alpha''_k(F_p)$. Let $\alpha_k'''(F_p)$ denotes the number of singularities with multiplicity kn or $kn + 1$ over p involved in $D^t(p)$. Then, we have $\alpha_k''(F_p) = \sum_{t=1}^{\eta_p} \alpha_k'''(F_p)$ by the definition of the decomposition $\tilde{R}_v(p) = D^1(p) + \cdots + D^{\eta_p}(p)$. Let $\alpha_k^{n\mathbb{Z}}(F_p)$, $\alpha_k^{n\mathbb{Z}+1}(F_p)$ respectively denote the number of singularities with multiplicity kn , $kn + 1$ over p . Similarly, we define $\alpha_k''^{n\mathbb{Z}}(F_p)$, $\alpha_k''^{n\mathbb{Z}+1}(F_p)$, etc.

Definition 1.7 (Singularity of type $(i \rightarrow i)$). Suppose that $n = 2$. If the exceptional curve E_x of the blow-up at a singularity x of R with multiplicity $2k + 1$ contains only one singularity y , then the multiplicity at y is $2k + 2$ and E_x contributes to $j_{0,1}''(F_p)$. Conversely, the exceptional curve E contributing to $j_{0,1}''(F_p)$ has such a pair (x, y) . Then we call the pair (x, y) a singularity of type $(2k + 1 \rightarrow 2k + 1)$ (cf. [18]). Let $\alpha_{(2k+1 \rightarrow 2k+1)}(F_p)$ be the number of singularities of type $(2k + 1 \rightarrow 2k + 1)$ over p (i.e., $s_{2k+1}(F_p)$ in the notation of [18]). Then we have

$$(1.12) \quad j_{0,1}''(F_p) = \sum_{k \geq 1} \alpha_{(2k+1 \rightarrow 2k+1)}(F_p).$$

We decompose

$$\alpha_{(2k+1 \rightarrow 2k+1)}(F_p) = \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{tr}}(F_p) + \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co}}(F_p)$$

and

$$\alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co}}(F_p) = \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},0}(F_p) + \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},1}(F_p)$$

as follows. Let $\alpha_{(2k+1 \rightarrow 2k+1)}^{\text{tr}}(F_p)$ be the number of singularities of type $(2k + 1 \rightarrow 2k + 1)$ over p at which any local branch of R_h intersects the fiber over p transversely. Let $\alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},0}(F_p)$ (resp. $\alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},1}(F_p)$) be the number of singularities (x, y) of type $(2k + 1 \rightarrow 2k + 1)$ over p such that the proper transform of the vertical component passing through x also passes through y and is not contained in R (resp. is contained in R).

Notation 1.8. For a condition or a Roman numeral $\mathcal{P}$, we put $\delta_{\mathcal{P}} = 1$ if the condition $\mathcal{P}$ holds or Γ_p is a singular fiber of type $\mathcal{P}$, and $\delta_{\mathcal{P}} = 0$ otherwise.

Let $C = C^{t,k}$ and assume that it is smooth. If C is on W_i , we drop the index and set $R = R_i$ for simplicity. Let $R' = R - C$. Let $x_1, \dots, x_l$ be all the points of $C \cap R'$. We put $x_{i,1} = x_i$ and $m_{i,1} = m_i$. We define $\psi_{i,1}: W_{i,1} \rightarrow W$ to be the blow-up at $x_{i,1}$ and put $E_{i,1} = \psi_{i,1}^{-1}(x_{i,1})$ and $R_{i,1} = \psi_{i,1}^* R - n[m_{i,1}/n]E_{i,1}$. Inductively, we define $x_{i,j}$, $m_{i,j}$ to be the intersection point of the proper transform of C and $E_{i,j-1}$, the multiplicity of $R_{i,j-1}$ at $x_{i,j}$, and if $m_{i,j} > 1$, we define $\psi_{i,j}: W_{i,j} \rightarrow W_{i,j-1}$, $E_{i,j}$ and $R_{i,j}$ to be the blow-up at $x_{i,j}$, the exceptional curve for $\psi_{i,j}$ and $R_{i,j} = \psi_{i,j}^* R_{i,j-1} - n[m_{i,j}/n]E_{i,j}$, respectively. Put $i_{\text{bm}} = \max\{j \mid m_{i,j} > 1\}$, that is, the number of blowing-ups occurring over x_i . We may assume that $i_{\text{bm}} \geq (i + 1)_{\text{bm}}$ for $i = 1, \dots, l - 1$ after rearranging the index if necessary. Put $t = R'C$ and $c = \sum_{i=1}^l i_{\text{bm}}$. If C is a fiber Γ of φ , t is the number of branch points r . If C is an exceptional curve, t is the multiplicity of R at the point to which C is contracted. Clearly, c is the number of blow-ups on C . Set $d_{i,j} = [m_{i,j}/n]$. Then the following lemmas hold (cf. [7]).

Lemma 1.9 ([7]). *We have*

$$(1.13) \quad \frac{t+c}{n} = \sum_{i=1}^l \sum_{j=1}^{i_{\text{bm}}} d_{i,j}.$$

This is a special case of the following lemma:

Lemma 1.10. *Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type (g, h, n) . Let C be a curve contained in R , L the proper transform of C on $\widetilde{W}$ and $x_1, x_2, \dots, x_c$ all the singularities of R on C (including infinitely near ones). Put $m_i = \text{mult}_{x_i}(R)$, $k_i = \text{mult}_{x_i}(C)$ and $d_i = \lfloor m_i/n \rfloor$. Then, we have*

$$\frac{RC - L^2}{n} = \sum_{i=1}^c k_i d_i.$$

Proof. We may assume that ψ_i is the blow-up at x_i for $i = 1, \dots, c$. Then, we can write $L = \tilde{\psi}^* C - \sum_{i=1}^c k_i \mathbf{E}_i$ and $\tilde{R} = \tilde{\psi}^* R - \sum_{i=1}^c n d_i \mathbf{E}_i$. Thus, we have $\tilde{R}L = RC - \sum_{i=1}^c n k_i d_i$. On the other hand, since $\tilde{R} - L$ and L are disjoint, we have $L^2 = \tilde{R}L$. From these equalities, the assertion follows. $\square$

Lemma 1.11 ([7]). *The following hold.*

- (1) *When $n \geq 3$, then $m_{i,j} \geq m_{i,j+1}$. When $n = 2$, then $m_{i,j} + 1 \geq m_{i,j+1}$ with equality holds only if $m_{i,j-1} \in 2\mathbb{Z}$ (if $j > 1$) and $m_{i,j} \in 2\mathbb{Z} + 1$.*
- (2) *If $m_{i,j-1} \in n\mathbb{Z} + 1$ and $m_{i,j} \in n\mathbb{Z}$, then $m_{i,j} > m_{i,j+1}$.*
- (3) *$m_{i,i_{\text{bm}}} \in n\mathbb{Z}$.*

Definition 1.12. By using the datum $\{m_{i,j}\}$, one can construct a diagram as in Table 1. We call it the *singularity diagram* of C .

TABLE 1. singularity diagram

#		
$(x_{1,1_{\text{bm}}}, m_{1,1_{\text{bm}}})$		
$\dots$		#
	$\dots$	$(x_{l,l_{\text{bm}}}, m_{l,l_{\text{bm}}})$
		$\dots$
$(x_{1,1}, m_{1,1})$	$\dots$	$(x_{l,1}, m_{l,1})$

On the top of the i -th column (indicated by # in Table 1), we write $\# = (i_{\text{max}} - i_{\text{bm}})$ if $i_{\text{bm}} < i_{\text{max}}$ and leave it blank when $i_{\text{bm}} = i_{\text{max}}$. We say that the singularity diagram of C is of type 0 (resp. of type 1) if $C \not\subset R$ (resp. $C \subset R$).

Definition 1.13. Let $\mathcal{D}^{t,k}$ be the singularity diagram of $C^{t,k}$. We call $\mathcal{D}^{t,1}, \mathcal{D}^{t,2}, \dots, \mathcal{D}^{t,j^{t(F_p)}}$ a sequence of singularity diagrams associated with $D^t(p)$.

Then the following lemma is clear.

Lemma 1.14. Let $\mathcal{D}^{t,1}, \mathcal{D}^{t,2}, \dots, \mathcal{D}^{t,j^t(F_p)}$ be a sequence of singularity diagrams associated with $D^t(p)$. Let $l^{t,k} := \#(R' \cap C^{t,k})$ and $(x_{i,j}^{t,k}, m_{i,j}^{t,k})$, $i = 1, \dots, l^{t,k}$, $j = 1, \dots, i_{\text{bm}}$ denote entries of $\mathcal{D}^{t,k}$. Let $(x_{i,j}^{t,p}, m_{i,j}^{t,p})$ be a singularity on $C^{t,p}$ such that $m_{i,j}^{t,p} \in n\mathbb{Z} + 1$, and $m_{i,j-1}^{t,p} \in n\mathbb{Z}$ when $j > 1$. Let $q > p$ be the integer such that $C^{t,q}$ is the exceptional curve for the blow-up at $x_{i,j}^{t,p}$. Then, for every $1 \leq p' \leq p$, i', j' satisfying $(x_{i',j'}^{t,p'}, m_{i',j'}^{t,p'}) = (x_{i,j}^{t,p}, m_{i,j}^{t,p})$, the diagram $\mathcal{D}^{t,q}$ has $(x_{i',j'+1}^{t,p'}, m_{i',j'+1}^{t,p'})$ as an entry in the bottom row.

Example 1.15. Suppose that t contributes to $\bar{\eta}_p$. Then $C^{t,k} = C'^{t,k}$ is a (-1) -curve and blown up $n - 1$ times for any k .

(1) If $n = 2$, then the point to which $C^{t,1}$ is contracted is a singularity of type $(m \rightarrow m)$ for some odd integer m . Indeed, $R'C^{t,1} = m$ and from Lemma 1.9, the singularity diagram of $C^{t,1}$ is the following:

$$\begin{array}{c} \boxed{m+1} \\ \mathcal{D}^{t,1} \end{array}$$

where we drop the symbol indicating the singular point on $C^{t,1}$ for simplicity. Since $m+1$ is even, we have $j^t(F_p) = 1$. This observation gives us

$$(1.14) \quad \bar{\eta}_p = \sum_{k \geq 1} \left(\alpha_{(2k+1 \rightarrow 2k+1)}^{\text{tr}}(F_p) + \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},0}(F_p) \right).$$

(2) Suppose that $n = 3$. Let m be the multiplicity of the singular point to which $C^{t,1}$ is contracted. Then $R'C^{t,1} = m$ and from Lemma 1.9, all possible singularity diagram of $C^{t,1}$ are the following:

$$\begin{array}{ccc} \text{(i)} \quad \begin{array}{|c|c|} \hline n_1 & n_2 \\ \hline \end{array} & \text{(ii)} \quad \begin{array}{|c|} \hline n_2 \\ \hline n_1 \\ \hline \end{array} & \text{(iii)} \quad \begin{array}{|c|} \hline n_1 \\ \hline m_1 \\ \hline \end{array} \\ \mathcal{D}^{t,1} & \mathcal{D}^{t,1} & \mathcal{D}^{t,1} \end{array}$$

where the integers $n_i \in 3\mathbb{Z}$ and $m_i \in 3\mathbb{Z} + 1$ satisfy that $m + 2 = n_1 + n_2$ in the case (i), $m + 2 = n_1 + n_2$ and $n_2 \leq n_1$ in the case (ii), $m + 3 = m_1 + n_1$ and $n_1 < m_1$ in the case (iii). If the diagram $\mathcal{D}^{t,1}$ is (i) or (ii), then $j^t(F_p) = 1$ since there are no $3\mathbb{Z} + 1$ type singularities on $C^{t,1}$. If the diagram $\mathcal{D}^{t,1}$ is (iii), then $j^t(F_p) > 1$ and the singularity diagram $\mathcal{D}^{t,2}$ of $C^{t,2}$ which is obtained by the blow-up at the singularity with multiplicity m_1 is (i) or (ii) from Lemma 1.14. Thus we have $j^t(F_p) = 2$.

(3) When $n \geq 4$, then the number $j^t(F_p)$ is not bounded. For example, we can consider the following sequence of singularity diagrams associated with $D^t(p)$ when $n = 4$:

$$\begin{array}{ccc}
\begin{array}{|c|c|} \hline n_1 & \\ \hline m_1 & n_2 \\ \hline \end{array} & \begin{array}{|c|c|} \hline n_3 & \\ \hline m_2 & n_4 \\ \hline \end{array} & \dots & \begin{array}{|c|c|} \hline n_{2N-1} & \\ \hline m_N & n_{2N} \\ \hline \end{array} \\
\mathcal{D}^{t,1} & \mathcal{D}^{t,2} & & \mathcal{D}^{t,N}
\end{array}$$

where $n_k \in 4\mathbb{Z}$, $m_k \in 4\mathbb{Z} + 1$ and $C^{t,k}$, $k \geq 2$ is the exceptional curve obtained by the blow-up at the multiplicity on $C^{t,k-1}$ with multiplicity m_{k-1} . From Lemma 1.9, we have $m_k + 4 = m_{k+1} + n_{2k+1} + n_{2k+2}$ for any $k \geq 1$.

Recall that the gonality $\text{gon}(C)$ of a non-singular projective curve C is the minimum of the degree of morphisms onto $\mathbb{P}^1$. The gonality $\text{gon}(f)$ of a fibered surface $f: S \rightarrow B$ is defined to be that of a general fiber (cf. [14]).

Proposition 1.16. *Let $\theta: F \rightarrow \Gamma$ be a totally ramified covering of degree n between smooth projective curves branched over r points. If $r \geq 2n \text{gon}(\Gamma)$, then $\text{gon}(F) = n \text{gon}(\Gamma)$. In particular, the gonality of a primitive cyclic covering fibration of type (g, h, n) is $n \text{gon}(\varphi)$, when $r \geq 2n \text{gon}(\varphi)$.*

Proof. Assume contrary that F has a morphism onto $\mathbb{P}^1$ of degree $k < n \text{gon}(\Gamma)$. This together with the covering $\theta: F \rightarrow \Gamma$ defines a morphism $\Phi: F \rightarrow \mathbb{P}^1 \times \Gamma$. If Φ is of degree m onto the image $\Phi(F)$, then m is a common divisor of n, k and the arithmetic genus of $\Phi(F)$ is $(n/m - 1)k/m + (h - 1)n/m + 1$ by the genus formula. Now, let F' be the normalization of $\Phi(F)$. Since the covering $F \rightarrow \Gamma$ factors through F' , we see that the induced covering $F' \rightarrow \Gamma$ of degree n/m is a totally ramified covering branched over r points. Then, by the Hurwitz formula, we have $2g(F') - 2 = (2h - 2)n/m + (n/m - 1)r$. Since the genus $g(F')$ of F' is not bigger than the arithmetic genus of $\Phi(F)$, we get $r \leq 2(k/m)$ when $n > m$, which is impossible, since $r \geq 2n \text{gon}(\Gamma)$ and $k < n \text{gon}(\Gamma)$. Thus, we get $n = m$. Then F' is isomorphic to Γ and therefore the morphism $F \rightarrow \mathbb{P}^1$ factors through Γ . Hence we have $k \geq n \text{gon}(\Gamma)$ by the definition of the gonality of Γ , which contradicts $k < n \text{gon}(\Gamma)$. A more careful study shows that any gonality pencil of F is the pull-back of a gonality pencil of Γ when $r > 2n \text{gon}(\Gamma)$. $\square$

2. PRIMITIVE CYCLIC COVERING FIBRATIONS OF TYPE $(g, 1, n)$

Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(g, 1, n)$. Since $\varphi: W \rightarrow B$ is a relatively minimal elliptic surface, K_φ is numerically equivalent to $\left(\chi_\varphi + \sum_{p \in B} \left(1 - \frac{1}{m_p}\right)\right) \Gamma$ by the canonical bundle formula, where m_p denotes the multiplicity of the fiber Γ_p of φ over p . In particular, we have $K_\varphi^2 = 0$. For $p \in B$, we put $\nu(F_p) = 1 - 1/m_p$ and $\nu = \sum_{p \in B} \nu(F_p)$. Then, we have $K_\varphi R = (\chi_\varphi + \nu)r$. Combining these equalities with (1.8), (1.9), (1.10) and (1.11), we get the following lemma:

Lemma 2.1. *The following equalities hold.*

$$\begin{aligned}
K_f^2 &= \sum_{k \geq 1} ((n+1)(n-1)k - n) \alpha_k + \frac{(n-1)^2}{n} (\alpha_0 - 2\varepsilon) \\
&\quad + \frac{(n+1)(n-1)r}{n} (\chi_\varphi + \nu) + \varepsilon. \\
\chi_f &= \frac{1}{12} (n-1)(n+1) \sum_{k \geq 1} k \alpha_k + \frac{(n-1)(2n-1)}{12n} (\alpha_0 - 2\varepsilon) \\
&\quad + \frac{(n+1)(n-1)r}{12n} (\chi_\varphi + \nu) + n\chi_\varphi. \\
e_f &= (n-1)\alpha_0 + n \sum_{k \geq 1} \alpha_k - (2n-1)\varepsilon + 12n\chi_\varphi.
\end{aligned}$$

For $p \in B$, we put $\chi_\varphi(F_p) = e_\varphi(\Gamma_p)/12$ and

$$\begin{aligned}
K_f^2(F_p) &= \sum_{k \geq 1} ((n+1)(n-1)k - n) \alpha_k(F_p) + \frac{(n-1)^2}{n} (\alpha_0(F_p) - 2\varepsilon(F_p)) \\
&\quad + \frac{(n+1)(n-1)r}{n} (\chi_\varphi(F_p) + \nu(F_p)) + \varepsilon(F_p), \\
\chi_f(F_p) &= \frac{1}{12} (n-1)(n+1) \sum_{k \geq 1} k \alpha_k(F_p) + \frac{(n-1)(2n-1)}{12n} (\alpha_0(F_p) - 2\varepsilon(F_p)) \\
&\quad + \frac{(n+1)(n-1)r}{12n} (\chi_\varphi(F_p) + \nu(F_p)) + n\chi_\varphi(F_p), \\
e_f(F_p) &= (n-1)\alpha_0(F_p) + n \sum_{k \geq 1} \alpha_k(F_p) - (2n-1)\varepsilon(F_p) + 12n\chi_\varphi(F_p).
\end{aligned}$$

Then, the following slope equality holds:

Theorem 2.2. *Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(g, 1, n)$. Then*

$$K_f^2 = \lambda_{g,1,n} \chi_f + \sum_{p \in B} \text{Ind}(F_p),$$

where $\lambda_{g,1,n} := 12(n-1)/(2n-1)$ and $\text{Ind}(F_p)$ is defined by

$$\begin{aligned}
\text{Ind}(F_p) &= n \sum_{k \geq 1} \left(\frac{(n+1)(n-1)}{2n-1} k - 1 \right) \alpha_k(F_p) + \frac{n-1}{2n-1} ((n+1)r - 12n) \chi_\varphi(F_p) \\
&\quad + \frac{(n+1)(n-1)r}{2n-1} \nu(F_p) + \varepsilon(F_p).
\end{aligned}$$

Moreover, if $r \geq \frac{12n}{n+1}$, then $\text{Ind}(F_p)$ is non-negative for any $p \in B$.

Proof. Since $K_f^2 = \sum_{p \in B} K_f^2(F_p)$, $\chi_f = \sum_{p \in B} \chi_f(F_p)$ and $K_f^2(F_p) - \lambda_{g,1,n} \chi_f(F_p) = \text{Ind}(F_p)$, the claim follows. $\square$

For an oriented compact real 4-dimensional manifold X , the signature $\text{Sign}(X)$ of X is defined to be the number of positive eigenvalues minus the number of negative eigenvalues of the intersection form on $H^2(X)$. From Lemma 2.1, we observe the local concentration of $\text{Sign}(S)$ to a finite number of fiber germs.

Corollary 2.3 (cf. [1]). *Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(g, 1, n)$. Then*

$$\text{Sign}(S) = \sum_{p \in B} \sigma(F_p),$$

where $\sigma(F_p)$ is defined by

$$\begin{aligned} \sigma(F_p) = & n \sum_{k \geq 1} \left(\frac{(n+1)(n-1)}{3} k - 1 \right) \alpha_k(F_p) + \left(\frac{(n-1)(n+1)r}{3n} - 8n \right) \chi_\varphi(F_p) \\ & + \frac{(n+1)(2n-1)}{3n} \varepsilon(F_p) + \frac{(n+1)(n-1)r}{3n} \nu(F_p) - \frac{(n+1)(n-1)}{3n} \alpha_0(F_p). \end{aligned}$$

Proof. By the index theorem (cf. [9, p. 126]), we have

$$\text{Sign}(S) = \sum_{p+q \equiv 0 \pmod{2}} h^{p,q}(S) = K_f^2 - 8\chi_f.$$

On the other hand, we can see that

$$\sigma(F_p) = K_f^2(F_p) - 8\chi_f(F_p)$$

by a computation. $\square$

3. UPPER BOUND OF THE SLOPE: THE CASE OF TYPE $(g, 1, n)$

In this section, we prove the following theorem:

Theorem 3.1. *Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(g, 1, n)$.*

(1) *If $n \geq 4$, then we have*

$$K_f^2 \leq \left(12 - \frac{12n^2}{r(n-1)(n+1)} \right) \chi_f.$$

(2) *If $n = 3$ and $g \geq 7$, then we have*

$$K_f^2 \leq \left(12 - \frac{24}{4g-17} \right) \chi_f.$$

(3) If $n = 2$ and $g \geq 3$, then we have

$$K_f^2 \leq \left(12 - \frac{2}{g-2}\right) \chi_f.$$

Corollary 3.2. *Let $f: S \rightarrow B$ be a relatively minimal bielliptic fibered surface of genus $g \geq 6$. Then, we have*


$$K_f^2 \leq \left(12 - \frac{2}{g-2}\right) \chi_f.$$

Proof. Let $f: S \rightarrow B$ be a relatively minimal fibered surface of genus g whose general fiber F is a double cover of a smooth curve Γ of genus h . If $g > 4h + 1$, an involution of the general fiber F of f over Γ is unique. Then, the fibration f has a global involution since it is relatively minimal (cf. [6]). Hence f is a primitive cyclic covering fibration of type $(g, h, 2)$. In particular, a relatively minimal bielliptic fibered surface of genus $g \geq 6$ is a primitive cyclic covering fibration of type $(g, 1, 2)$. $\square$


Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(g, 1, n)$. We fix $p \in B$. Let $m = m_p$ be the multiplicity of the fiber Γ_p of φ over p . Since $h = 1$, we have $j_{b,\bullet}(F_p) = 0$ for any $b \geq 2$. From the classification of singular fibers of relatively minimal elliptic surfaces ([12]), we have the following lemma for u -vertical type singularities:

Lemma 3.3. *There exist no u -vertical type singularities of R for $u \geq 4$. All possible 3-vertical type singularities are as follows.*


Type (II) : Γ_p is a singular fiber of type (II) in the Kodaira's table ([12])(i.e., it is a singular rational curve with one cusp) and it is contained in R . The cusp on Γ_p is a singularity of type $n\mathbb{Z} + 1$ and the singularity at which the proper transform of Γ_p and the exceptional curve E_1 for the blow-up at the cusp intersect is also of type $n\mathbb{Z} + 1$. Then, the proper transforms of Γ_p and E_1 and the exceptional curve E_2 for the blow-up at this singularity form a 3-vertical type singularity.



Type (III) : Γ_p is a singular fiber of type (III) in the Kodaira's table (i.e., it consists of two nonsingular rational curves intersecting each other at one point of order two) and it is contained in R . The singularity on Γ_p is a singularity of type $n\mathbb{Z} + 1$. Then, the proper transforms of Γ_p and the exceptional curve E_1 for the blow-up at this singularity form a 3-vertical type singularity.



Type (IV) : Γ_p is a singular fiber of type (IV) in the Kodaira's table (i.e. it consists of three nonsingular rational curves intersecting one another at one point transversely) and it is contained in R . The singularity on Γ_p is a 3-vertical type singularity.



In particular, we have $\iota^{(u)}(F_p) = \kappa^{(u)}(F_p) = 0$ for $u \geq 4$ and $0 \leq \iota^{(3)}(F_p) + \kappa^{(3)}(F_p) \leq 1$.

Next, we give a lower bound of $\alpha_0^+(F_p)$ by using $\iota(F_p)$ and $\kappa(F_p)$.

Lemma 3.4. *We have*

$$\alpha_0^+(F_p) \geq \left(1 - \frac{1}{m}\right) r + (n-2)(\iota(F_p) + 2\kappa(F_p)) + \beta_p.$$

where $\beta_p := \delta_{n \neq 2}((n-7)\delta_{\text{II}} - (n+1)\delta_{\text{III}} - 2\delta_{\text{IV}})\iota^{(3)}(F_p) + \delta_{n=2} \sum_{k \geq 1} 2k\alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co}}(F_p)$.

Proof. Let $\tilde{\Gamma}_p = m\tilde{D}_p$ and $\tilde{D}_p = \sum_i m_i G_i$ the irreducible decomposition. Then we have

$$\begin{aligned} \alpha_0^+(F_p) &= r - \#(\text{Supp}(\tilde{R}_h) \cap \text{Supp}(\tilde{\Gamma}_p)) \\ &\geq r - \sum_i \tilde{R}_h G_i \\ &= \left(1 - \frac{1}{m}\right) r + \sum_i (m_i - 1) \tilde{R}_h G_i. \end{aligned}$$

For a $(t, 2)$ or $(t, 3)$ -vertical $n\mathbb{Z}$ type singularity x , we denote by E_x^t the exceptional curve for the blow-up at x . Let m_x^t be the multiplicity of $\tilde{D}_p$ along $\hat{E}_x^t$, the proper transform of E_x^t on $\widetilde{W}$. Then, we have

$$\sum_i (m_i - 1) \tilde{R}_h G_i \geq \sum_{t=1}^{\eta_p} \sum_{x: (t, u) \ n\mathbb{Z}, u \geq 2} (m_x^t - 1) \tilde{R}_h \hat{E}_x^t.$$

If there exists a singular point of type $n\mathbb{Z}$ on E_x^t , we replace E_x^t with the exceptional curve E obtained by blowing up at this point. Repeating this procedure, we may assume that there exist no singular points of type $n\mathbb{Z}$ on E_x^t . If there exists a singular point of type $n\mathbb{Z} + 1$ on E_x^t , the proper transform of the exceptional curve obtained by blowing up at this point belongs to other $D^u(p)$. Since the multiplicity of $\tilde{\Gamma}_p$ along it is not less than $m_x^t > 1$, we do not have to consider this situation. Thus, we may assume that there exist no singular points

on E_x^t and we have $\widetilde{R}_h \widehat{E}_x^t \geq n - 2$ if x is a 2-vertical type singularity, and $\widetilde{R}_h \widehat{E}_x^t \geq n - 3$ if x is a 3-vertical type singularity. We can see that $\sum_{x:(t,2) \ n\mathbb{Z}} m_x^t \geq 2\iota^t(F_p) + 2\kappa^t(F_p)$ for any t with $\iota^{t,(3)}(F_p) = 0$. Thus, if $\iota^{(3)}(F_p) = 0$, we have

$$\alpha_0^+(F_p) \geq \left(1 - \frac{1}{m}\right) r + (n - 2)(\iota(F_p) + 2\kappa(F_p)).$$

If $\iota^{(3)}(F_p) = 1$, then $m = 1$ and Γ_p is of type (II), (III) or (IV) from Lemma 3.3. We may assume that $D^1(p) \neq 0$. Let x_0 be the 3-vertical $n\mathbb{Z}$ type singularity over p . Suppose that Γ_p is of type (II). Then, we can see that $m_{x_0}^1 = 6$ and $\sum_{x:(1,2) \ n\mathbb{Z}} m_x^1 \geq 2\iota^{1,(2)}(F_p) + 2(\kappa^1(F_p) - 1)$. Then we have

$$\begin{aligned} \alpha_0^+(F_p) &\geq \left(1 - \frac{1}{m}\right) r + 5(n - 3) + (n - 2)(\iota^{1,(2)}(F_p) + 2(\kappa^1(F_p) - 1)) \\ &\quad + (n - 2) \sum_{t=2}^{\eta_p} (\iota^t(F_p) + 2\kappa^t(F_p)) \\ &= \left(1 - \frac{1}{m}\right) r + n - 7 + (n - 2)(\iota(F_p) + 2\kappa(F_p)). \end{aligned}$$

Suppose that Γ_p is of type (III). Then, we can see that $m_{x_0}^1 = 4$ and $\sum_{x:(1,2) \ n\mathbb{Z}} m_x^1 \geq 2\iota^{1,(2)}(F_p) + 2(\kappa^1(F_p) - 1)$. Then we have

$$\begin{aligned} \alpha_0^+(F_p) &\geq \left(1 - \frac{1}{m}\right) r + 3(n - 3) + (n - 2)(\iota^{1,(2)}(F_p) + 2(\kappa^1(F_p) - 1)) \\ &\quad + (n - 2) \sum_{t=2}^{\eta_p} (\iota^t(F_p) + 2\kappa^t(F_p)) \\ &= \left(1 - \frac{1}{m}\right) r - n - 1 + (n - 2)(\iota(F_p) + 2\kappa(F_p)). \end{aligned}$$

Suppose that Γ_p is of type (IV). Then, we can see that $m_{x_0}^1 = 3$ and $\sum_{x:(1,2) \ n\mathbb{Z}} m_x^1 \geq 2\iota^{1,(2)}(F_p) + 2\kappa^1(F_p)$. Then we have

$$\begin{aligned} \alpha_0^+(F_p) &\geq \left(1 - \frac{1}{m}\right) r + 2(n - 3) + (n - 2)(\iota^{1,(2)}(F_p) + 2\kappa^1(F_p)) \\ &\quad + (n - 2) \sum_{t=2}^{\eta_p} (\iota^t(F_p) + 2\kappa^t(F_p)) \\ &= \left(1 - \frac{1}{m}\right) r - 2 + (n - 2)(\iota(F_p) + 2\kappa(F_p)). \end{aligned}$$

Suppose that $n = 2$. For a $(2k + 1 \rightarrow 2k + 1)$ singularity (x, y) , let E_y denotes the exceptional curve for the blow-up at y and m_y the multiplicity of $\widetilde{\Gamma}_p$ along $\widehat{E}_y$, the proper transform of

E_y . Then we have

$$\sum_i (m_i - 1) \tilde{R}_h G_i \geq \sum_{k \geq 1} \sum_{(x,y):(2k+1 \rightarrow 2k+1)} (m_y - 1) \tilde{R}_h \hat{E}_y.$$

By an argument similar to the above, we may assume that there are no singular points on E_y . Then we have $\tilde{R}_h \hat{E}_y = 2k$ for any $(2k+1 \rightarrow 2k+1)$ singularity (x, y) . On the other hand, we have $m_y \geq 2$ for any $(2k+1 \rightarrow 2k+1)$ singularity (x, y) involved in $\alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co}}(F_p)$. Thus, we obtain $\sum_{(x,y):(2k+1 \rightarrow 2k+1)} (m_y - 1) \tilde{R}_h \hat{E}_y \geq 2k \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co}}(F_p)$. $\square$

We can translate the index α'' into other indices as follows.

Lemma 3.5. *The following equalities hold.*

$$(3.1) \quad \sum_{k \geq 1} \alpha_k''(F_p) = \eta_p'' + \sum_{a \geq 1} a n j_{1,a}'(F_p) + \sum_{a \geq 1} (a n - 2 - \delta_{m_{I_1, \Pi}}) j_{0,a}'(F_p) \\ + \sum_{a \geq 1} (a n - 1) j_{0,a}''(F_p) - \iota(F_p) - \kappa(F_p).$$

$$(3.2) \quad \sum_{k \geq 1} k \alpha_k''(F_p) = \gamma_p + \sum_{a \geq 1} a j_{\bullet, a}(F_p) + \sum_{k \geq 1} k (\alpha_k^{n\mathbb{Z}+1}(F_p) - \iota_k(F_p) - \kappa_k(F_p)).$$

where $\gamma_p = \sum_{t=1}^{\eta_p} \gamma_p^t$ and γ_p^t is defined to be the following (1), (2), (3):

- (1) $\gamma_p^t = d^{t,1}$ if $D^t(p) = 0$, where $m^{t,1}$ is the multiplicity of the singularity to which $C^{t,1}$ contracts and $d^{t,1} = [m^{t,1}/n]$.
- (2) $\gamma_p^t = \sum_{k=1}^{j^{tt}(F_p)} R C^{tt,k} / n$ if $D^t(p) \neq 0$ and any $C^{tt,k}$ is smooth.
- (3) $\gamma_p^t = r/mn - d^{t,1}$ if $C^{tt,1} = (\Gamma_p)_{\text{red}}$ is singular, where $m^{t,1}$ is the multiplicity of the singular point of R which is singular for $C^{tt,1}$ and $d^{t,1} = [m^{t,1}/n]$.

Proof. Let $a^{t,k}, a^{tt,k}$ be the integers such that $(L^{t,k})^2 = -a^{t,k}n$, $(L^{tt,k})^2 = -a^{tt,k}n$. If $C^{t,k}$ is smooth, then $C^{tt,k}$ is blown up $a^{t,k}n + (C^{t,k})^2$ times. If $C^{t,k}$ is a singular rational curve, then $C^{tt,k}$ is blown up $a^{t,k}n - 3$ times. Since $C^{tt,k}$ is a (-1) -curve, $C^{ttt,k}$ is blown up $a^{tt,k}n - 1$ times. Hence, if $D^t(p) \neq 0$ and every $C^{t,k}$ is smooth (resp. $C^{tt,1}$ is singular rational), the number of singular points associated with $D^t(p)$ is $\sum_k (a^{t,k}n + (C^{t,k})^2) +$ (resp. $\sum_k (a^{t,k}n - 3) +$) $\sum_k (a^{tt,k}n - 1) - \iota^t(F_p) - \kappa^t(F_p)$. Namely, we have

$$\sum_{k \geq 1} \alpha_k^{tt}(F_p) = \sum_{a \geq 1} a n j_{1,a}^{tt}(F_p) + \sum_{a \geq 1} (a n - 2 - \delta_{m_{I_1, \Pi}}) j_{0,a}^{tt}(F_p) + \sum_{a \geq 1} (a n - 1) j_{0,a}^{ttt}(F_p) - \iota^t(F_p) - \kappa^t(F_p).$$

If $D^t(p) = 0$, we have

$$\sum_{k \geq 1} \alpha_k^{tt}(F_p) = 1 + \sum_{a \geq 1} (a n - 1) j_{0,a}^{ttt}(F_p) - \iota^t(F_p) - \kappa^t(F_p).$$

Summing up for $t = 1, \dots, \eta_p$, we have (3.1).

Let $r^{t,k} = RC'^{t,k}$, $m^{t,k}$ the multiplicity of R at the point to which $C''^{t,k}$ is contracted and $d^{t,k} = [m^{t,k}/n]$. Let $x_1^{t,k}, \dots, x_{c'}^{t,k}$ (resp. $x_1''^{t,k}, \dots, x_{c''}''^{t,k}$) be all the singular points on $C'^{t,k}$ (resp. on $C''^{t,k}$), including infinitely near ones. Put $m_i^{t,k} = \text{mult}_{x_i^{t,k}}(R)$, $d_i^{t,k} = [m_i^{t,k}/n]$, $m_i''^{t,k} = \text{mult}_{x_i''^{t,k}}(R)$ and $d_i''^{t,k} = [m_i''^{t,k}/n]$. Applying Lemma 1.10 to $C'^{t,k}$ and $C''^{t,k}$, we get that $r^{t,k}/n + a'^{t,k} = \sum_i d_i^{t,k}$ if $C'^{t,k}$ is smooth, $r^{t,1}/n + a'^{t,1} = d_1^{t,1} + \sum_i d_i^{t,1}$ if $C'^{t,1}$ is singular rational, and $d^{t,k} + a'^{t,k} = \sum_i d_i^{t,k}$. If $D^t(p) \neq 0$ and every $C'^{t,k}$ is smooth, then

$$\begin{aligned} \sum_k \frac{r^{t,k}}{n} + \sum_{a \geq 1} a j_{\bullet,a}^t(F_p) + \sum_{k \geq 1} k \alpha_k^{t,n\mathbb{Z}+1}(F_p) &= \sum_k \left(\frac{r^{t,k}}{n} + a'^{t,k} \right) + \sum_k (d^{t,k} + a''^{t,k}) \\ &= \sum_k \sum_i d_i^{t,k} + \sum_k \sum_i d_i''^{t,k} \\ &= \sum_{k \geq 1} k (\alpha_k''^t(F_p) + \iota_k^t(F_p) + \kappa_k^t(F_p)). \end{aligned}$$

Similarly, if $D^t(p) \neq 0$ and $C'^{t,1}$ is singular rational, we have

$$\frac{r}{nm} - d_1^{t,1} + \sum_{a \geq 1} a j_{\bullet,a}^t(F_p) + \sum_{k \geq 1} k \alpha_k^{t,n\mathbb{Z}+1}(F_p) = \sum_{k \geq 1} k (\alpha_k''^t(F_p) + \iota_k^t(F_p) + \kappa_k^t(F_p)).$$

If $D^t(p) = 0$, then

$$\begin{aligned} \sum_{a \geq 1} a j_{\bullet,a}^t(F_p) + \sum_{k \geq 1} k \alpha_k^{t,n\mathbb{Z}+1}(F_p) &= \sum_k (d^{t,k} + a''^{t,k}) \\ &= \sum_k \sum_i d_i''^{t,k} \\ &= \sum_{k \geq 1} k (\alpha_k''^t(F_p) + \iota_k^t(F_p) + \kappa_k^t(F_p)) - d^{t,1}. \end{aligned}$$

Summing up for $t = 1, \dots, \eta_p$, we get (3.2). □

Lemma 3.6. *The following hold.*

$$\begin{aligned} \gamma_p &\leq \left(\frac{r}{n} - j'_{0,1}(F_p) \delta_{n=2} - \delta_{m\mathbf{I}_1, \mathbf{II}} \right) \delta_{\eta'_p \neq 0} + \left(\frac{r}{n} - 1 \right) \eta_p'' \\ \iota(F_p) &= j(F_p) - \eta_p + \delta_{\text{cyc}}, \end{aligned}$$

where δ_{cyc} is defined to be 1 if the following (1), (2), (3) and (4) hold and $\delta_{\text{cyc}} = 0$ otherwise.

(1) Γ_p is a singular fiber of type $(m\mathbf{I}_k)_{k \geq 1}$, $(\mathbf{II})$, $(\mathbf{III})$ or $(\mathbf{IV})$.

(2) Any irreducible component of Γ_p is contained in R .

(3) $\iota^{(3)}(F_p) = \kappa^{(3)}(F_p) = 0$.

(4) The multiplicity of the singular point of R which is singular for $(\Gamma_p)_{\text{red}}$ belongs to $n\mathbb{Z} + 1$ if Γ_p is a singular fiber of type $(m\mathbf{I}_1)$ or $(\mathbf{II})$.

Proof. By the definition of γ_p , the first inequality is clear. We consider the following graph $\mathbf{G}^t$: The vertex set $V(\mathbf{G}^t)$ is defined by the symbol set $\{v^{t,k}\}_{k=1}^{j^t(F_p)}$. The edge set $E(\mathbf{G}^t)$ is defined by the symbol set $\{e_x\}_x \cup \{e_y\}_y \cup \{e'_y\}_y$, where x, y respectively move among $(t, 2)$, $(t, 3)$ -vertical $n\mathbb{Z}$ type singularities. If the proper transform of $C^{t,k}$ meets that of $C^{t,k'}$ at a $(t, 2)$ -vertical $n\mathbb{Z}$ type singularity x , the edge e_x connects $v^{t,k}$ and $v^{t,k'}$. If the proper transforms of $C^{t,k}$, $C^{t,k'}$ and $C^{t,k''}$ ($k < k' < k''$) intersects in a $(t, 3)$ -vertical $n\mathbb{Z}$ type singularity y , the edge e_y connects $v^{t,k}$ and $v^{t,k'}$, and e'_y connects $v^{t,k'}$ and $v^{t,k''}$. By the definition of the decomposition $\tilde{R}_v(p) = D^1(p) + \cdots + D^{\eta_p}(p)$, the graph $\mathbf{G}^t$ is connected for any $t = 1, \dots, \eta_p$. Clearly, $\iota(F_p)$ is the cardinality of $E(\mathbf{G}^t)$. Thus, the number of cycles in $\mathbf{G}^t$ is $\iota(F_p) - j^t(F_p) + 1$. One sees that $\mathbf{G}^t$ has at most one cycle, and it has one cycle only if $\{C^{t,k}\}_k$ contains all irreducible components of Γ_p . Hence at most one $\mathbf{G}^t$ has one cycle. We can see that $\mathbf{G}^t$ has one cycle for some t if and only if $\delta_{\text{cyc}} = 1$. Thus, we get $\iota(F_p) = j(F_p) - \eta_p + \delta_{\text{cyc}}$. $\square$

For any singular point x of R , the multiplicity $\text{mult}_x(R)$ at x does not exceed $r/m + 1$ since $R(\Gamma_p)_{\text{red}} = r/m$. Thus we have $\alpha_k = 0$ for $k \geq r/nm + 1$. Moreover, the following lemma holds.

Lemma 3.7. *If $n \geq 3$, then we have $\alpha_{\frac{r}{nm}}^{n\mathbb{Z}+1}(F_p) = 0$. If $n = 2$, then we have $\kappa_{\frac{r}{2m}}(F_p) = 0$.*

Proof. If $\alpha_{\frac{r}{nm}}^{n\mathbb{Z}+1}(F_p) \neq 0$, then there exists an irreducible component C of Γ_p contained in R and a singular point x of R on C with multiplicity $r/m + 1$ such that any local horizontal branch of R around x is not tangential to C since $RD_p = r/m$. Then, the exceptional curve E for the blow-up at x and the proper transform of C form a singular point of multiplicity 2. Hence we have $n = 2$ from Lemma 1.2. It is clear that all singular points with multiplicity $r/m + 1$ are infinitely near to x and the exceptional curves for blow-ups of these singularities form a chain. In particular, any singular point with multiplicity $r/m + 1$ is a 1-vertical type singularity. $\square$

To prove Theorem 3.1, we need some inequalities among several indices.

Lemma 3.8. (1) *The following holds.*

$$\begin{aligned} \sum_{k \geq 1} k (\alpha_k^{n\mathbb{Z}+1}(F_p) - \kappa_k(F_p)) &\leq \left(\frac{r}{n} - 1\right) (j''(F_p) - \kappa(F_p)) \\ &\quad + \left(\frac{r}{n} - 2\right) \kappa^{(3)}(F_p) + \alpha_{\frac{r}{nm}}^{n\mathbb{Z}+1}(F_p). \end{aligned}$$

(2) *If $n = 2$, then the following holds more strongly.*

$$\begin{aligned} \sum_{k \geq 1} k (\alpha_k^{2\mathbb{Z}+1}(F_p) - \kappa_k(F_p)) &\leq \sum_{k \geq 1} k \alpha_{(2k+1 \rightarrow 2k+1)}(F_p) + \left(\frac{r}{2} - 1\right) \left(\sum_{a \geq 2} j''_{0,a}(F_p) - \kappa(F_p)\right) \\ &\quad + \left(\frac{r}{2} - 2\right) \kappa^{(3)}(F_p) + \alpha_{\frac{r}{2m}}^{2\mathbb{Z}+1}(F_p). \end{aligned}$$

Proof. From Lemma 3.7, we have

$$\begin{aligned} \sum_{k \geq 1} k (\alpha_k^{n\mathbb{Z}+1}(F_p) - \kappa_k(F_p)) &= \sum_{k=1}^{\frac{r}{nm}-1} k (\alpha_k^{n\mathbb{Z}+1}(F_p) - \kappa_k(F_p)) + \frac{r}{nm} \alpha_{\frac{r}{nm}}^{n\mathbb{Z}+1}(F_p) \\ &= \sum_{k=1}^{\frac{r}{nm}-1} k (\alpha_k^{n\mathbb{Z}+1}(F_p) - \kappa_k^{(2)}(F_p) - \kappa_k^{(3)}(F_p)) - \sum_{k=1}^{\frac{r}{nm}-1} k \kappa_k^{(3)}(F_p) + \frac{r}{nm} \alpha_{\frac{r}{nm}}^{n\mathbb{Z}+1}(F_p) \end{aligned}$$

Since $\alpha_k^{n\mathbb{Z}+1}(F_p) - \kappa_k^{(2)}(F_p) - \kappa_k^{(3)}(F_p) \geq 0$ and $\sum_{k=1}^{\frac{r}{nm}-1} k \kappa_k^{(3)}(F_p) = k_0 \kappa^{(3)}(F_p)$ for some $1 \leq k_0 \leq r/n - 1$ from Lemma 3.3, we have

$$\begin{aligned} &\sum_{k=1}^{\frac{r}{nm}-1} k (\alpha_k^{n\mathbb{Z}+1}(F_p) - \kappa_k^{(2)}(F_p) - \kappa_k^{(3)}(F_p)) - \sum_{k=1}^{\frac{r}{nm}-1} k \kappa_k^{(3)}(F_p) + \frac{r}{nm} \alpha_{\frac{r}{nm}}^{n\mathbb{Z}+1}(F_p) \\ &\leq \left(\frac{r}{n} - 1\right) \left(\sum_{k=1}^{\frac{r}{nm}-1} \alpha_k^{n\mathbb{Z}+1}(F_p) - \kappa^{(2)}(F_p) - \kappa^{(3)}(F_p) \right) - k_0 \kappa^{(3)}(F_p) + \frac{r}{n} \alpha_{\frac{r}{nm}}^{n\mathbb{Z}+1}(F_p) \\ &= \left(\frac{r}{n} - 1\right) \left(\sum_{k=1}^{\frac{r}{nm}-1} \alpha_k^{n\mathbb{Z}+1}(F_p) - \kappa(F_p) \right) + \left(\frac{r}{n} - 1 - k_0\right) \kappa^{(3)}(F_p) + \frac{r}{n} \alpha_{\frac{r}{nm}}^{n\mathbb{Z}+1}(F_p). \end{aligned}$$

Combining the above inequality with $j''(F_p) = \sum_{k=1}^{\frac{r}{nm}} \alpha_k^{n\mathbb{Z}+1}(F_p)$ and $r/n - 1 - k_0 \leq r/n - 2$, the assertion (1) follows.

Assume $n = 2$. Note that any $(2k+1 \rightarrow 2k+1)$ singularity is not involved in $\kappa(F_p)$. Then we have

$$\begin{aligned} \sum_{k \geq 1} k (\alpha_k^{2\mathbb{Z}+1}(F_p) - \kappa_k(F_p)) &= \sum_{k=1}^{\frac{r}{2m}-1} k (\alpha_k^{2\mathbb{Z}+1}(F_p) - \alpha_{(2k+1 \rightarrow 2k+1)}(F_p) - \kappa_k(F_p)) \\ &\quad + \sum_{k \geq 1} k \alpha_{(2k+1 \rightarrow 2k+1)}(F_p) + \frac{r}{2m} \alpha_{\frac{r}{2m}}^{2\mathbb{Z}+1}(F_p). \end{aligned}$$

Similarly as in (1), the assertion (2) follows. □

Lemma 3.9. *If $n = 2$, then we have*

$$\kappa(F_p) \leq \frac{2}{3} \sum_{a \geq 2} (a-1) j_{\bullet, a}(F_p) - \frac{2}{3} \alpha_{\frac{r}{2m}}^{2\mathbb{Z}+1}(F_p).$$

Proof. It is sufficient to show that

$$(3.3) \quad \kappa^t(F_p) \leq \frac{2}{3} \sum_{a \geq 2} (a-1) j_{0, a}^t(F_p) - \frac{2}{3} \alpha_{\frac{r}{2m}}^{t, 2\mathbb{Z}+1}(F_p)$$

for any t . If $\kappa^t(F_p) = 0$, then it is clear. Thus, we may assume $\kappa^t(F_p) > 0$. Then clearly we have

$$(3.4) \quad j^t(F_p) \geq \kappa^{t,(2)}(F_p) + \kappa^{t,(3)}(F_p) + \alpha_{\frac{r}{2m}}^{t,2\mathbb{Z}+1}(F_p) + 2.$$

Since any blow-up at a (t, u) -vertical type singularity contributes $-u$ to the number

$$\sum_{k \geq 1} (L^{t,k})^2 = - \sum_{a \geq 1} 2aj_{\bullet,a}^t(F_p)$$

and Γ_p contains no u -vertical type singularity for $u \geq 2$ if Γ_p is of type $({}_m\mathrm{I}_0)$, we get

$$\begin{aligned} \sum_{a \geq 1} 2aj_{\bullet,a}^t(F_p) &\geq j_{1,\bullet}^t(F_p) + (2 + \delta_{m\mathrm{I}_1,\mathrm{II}})j_{0,\bullet}^t(F_p) + j_{0,\bullet}^{\prime\prime t}(F_p) \\ &\quad + \alpha_{\frac{r}{2m}}^{t,2\mathbb{Z}+1}(F_p) + \sum_{u=2,3} u \left(\iota^{t,(u)}(F_p) + \kappa^{t,(u)}(F_p) \right). \end{aligned}$$

Combining this inequality with $\iota^t(F_p) \geq j^t(F_p) - 1$ and (3.4), we have

$$\begin{aligned} \sum_{a \geq 1} 2aj_{\bullet,a}^t(F_p) &\geq 3j^t(F_p) + (1 + \delta_{m\mathrm{I}_1,\mathrm{II}})j_{0,\bullet}^t(F_p) + \alpha_{\frac{r}{2m}}^{t,2\mathbb{Z}+1}(F_p) \\ &\quad + 2\kappa^t(F_p) - 2 - (\iota^{t,(3)}(F_p) + \kappa^{t,(3)}(F_p)) \\ &\geq 2j^t(F_p) + (1 + \delta_{m\mathrm{I}_1,\mathrm{II}})j_{0,\bullet}^t(F_p) + 2\alpha_{\frac{r}{2m}}^{t,2\mathbb{Z}+1}(F_p) \\ &\quad + 3\kappa^t(F_p) - \iota^{t,(3)}(F_p) - 2\kappa^{t,(3)}(F_p). \end{aligned}$$

On the other hand, it is easily seen that

$$(1 + \delta_{m\mathrm{I}_1,\mathrm{II}})j_{0,\bullet}^t(F_p) - \iota^{t,(3)}(F_p) - 2\kappa^{t,(3)}(F_p) \geq 0.$$

Hence we get (3.3), as desired. $\square$

Lemma 3.10. (1) *If $n = 3$, then the following hold.*

(1, i) *If $j_{0,1}^t(F_p) \leq 2$ for any t , then*

$$\frac{1}{2}j'_{0,1}(F_p) \leq \eta'_p - \delta_{\mathrm{cyc}}.$$

(1, ii) *If $j_{0,1}^t(F_p) = 3$ for some t , then Γ_p is a singular fiber of type (IV) , (I_k^*) , (II^*) , (III^*) or (IV^*) and*

$$\frac{1}{3}j'_{0,1}(F_p) \leq \eta'_p, \quad \delta_{\mathrm{cyc}} = 0.$$

(1, iii) *If $j_{0,1}^t(F_p) = 4$ for some t , then Γ_p is a singular fiber of type (I_k^*) and any component of Γ_p is contained in R . Moreover, we have $\eta'_p = 1$, $j'_{0,1}(F_p) = 4$ and $\delta_{\mathrm{cyc}} = 0$.*

(2) *If $n = 2$, then the following hold.*

(2, i) *If $j_{0,2,\mathrm{odd}}^t(F_p) \leq 2$ for any t , then*

$$j'_{0,1}(F_p) + \frac{1}{2}j'_{0,2,\mathrm{odd}}(F_p) \leq \eta'_p - \delta_{\mathrm{cyc}},$$

where $j'_{0,2,\text{odd}}(F_p)$ denotes the number of irreducible components C of Γ_p involved in $j'_{0,2}(F_p)$ which has a singular point of R of odd multiplicity.

(2, ii) If $j_{0,2,\text{odd}}^t(F_p) = 3$ for some t , then Γ_p is a singular fiber of type (IV), (I_k^*) , (II^*) , (III^*) or (IV^*) and

$$j'_{0,1}(F_p) + \frac{1}{3}j'_{0,2,\text{odd}}(F_p) \leq \eta'_p, \quad \delta_{\text{cyc}} = 0.$$

(2, iii) If $j_{0,2,\text{odd}}^t(F_p) = 4$ for some t , then Γ_p is a singular fiber of type (I_k^*) and any component of Γ_p is contained in R . Moreover, we have $\eta'_p = 1$, $j'_{0,1}(F_p) = 0$, $j'_{0,2,\text{odd}}(F_p) = 4$ and $\delta_{\text{cyc}} = 0$.

Proof. If $n = 3$, then any curve C in Γ_p contributing to $j'_{0,1}(F_p)$ intersects at most one component of Γ_p contained in R , since C is blown up just once. Thus, considering the classification of singular fibers of elliptic surfaces, we can show easily the assertion (1).

Suppose that $n = 2$. Any curve in Γ_p contributing to $j'_{0,1}(F_p)$ is not blown up and any curve in Γ_p contributing to $j'_{0,2,\text{odd}}(F_p)$ intersects at most one component of Γ_p contained in R . Hence we can show the assertion (2) similarly. $\square$

Lemma 3.11. (1) If $n = 3$, then we have

$$j''_{0,1}(F_p) \leq 2\eta''_p + \sum_{a \geq 1} 2aj'_{1,a}(F_p) + \sum_{a \geq 2} (2a - 2)j'_{0,a}(F_p) + \sum_{a \geq 2} (2a - 1)j''_{0,a}(F_p).$$

(2) If $n = 2$, then we have

$$\begin{aligned} \sum_{k \geq 1} \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},1}(F_p) &\leq j'_{0,2,\text{odd}}(F_p) + \sum_{a \geq 3} (a - 1)j'_{0,a}(F_p) + \sum_{a \geq 2} (a - 1)j'_{1,a}(F_p) \\ &\quad + \sum_{a \geq 3} (a - 2)j''_{0,a}(F_p) + \widehat{\eta}_p. \end{aligned}$$

Proof. Suppose that $n = 3$. Let $C_1, \dots, C_{j_{0,1}'''(F_p)}$ be all (-1) -curves in $\{C^{t,k}\}_k$ contributing to $j_{0,1}'''(F_p)$ and x_i the point to which C_i contracts for $i = 1, \dots, j_{0,1}'''(F_p)$. If $C_i \neq C^{t,1}$, then x_i is contained in some $C^{t,k}$. If $C^{t,k}$ contributes to $j_{0,1}'''(F_p)$ and $k = 1$, then $j_{0,1}'''(F_p) = j_{0,1}'''(F_p) = 2$ from Example 1.15 (2). If $C^{t,k}$ contributes to $j_{0,1}'''(F_p)$ and $k \neq 1$, then the point x_i to which $C^{t,k}$ contracts is contained in another $C^{t,k'}$ which does not contribute to $j_{0,1}'''(F_p)$ from the argument of Example 1.15 (2). Moreover, x_i is also contained in $C^{t,k'}$ since the singularity diagram of $C^{t,k}$ is type (iii) in Example 1.15 (2) and Lemma 1.9. For a curve $C^{t,k}$ which does not contribute to $j_{0,1}'''(F_p)$, we consider how many points among $x_1, \dots, x_{j_{0,1}'''(F_p)}$ it contains.

(i) Assume that $C^{t,k}$ contributes to $j_{0,a}'''(F_p)$ for some $a \geq 2$. Then $C^{t,k}$ is blown up $3a - 1$ times. Let $(x_{i,j}, m_{i,j})$, $i = 1, \dots, l$, $j = 1, \dots, i_{\text{bm}}$ be entries of the singularity diagram of $C^{t,k}$. We consider a subset of entries of the i -th column of its diagram $\{(x_{i,j}, m_{i,j})\}_{j=j_0+1, \dots, j_0+N}$ satisfying that

(*) $m_{i,j_0} \in 3\mathbb{Z}$ if $j_0 > 0$, $m_{i,j} \in 3\mathbb{Z} + 1$ for $j_0 < j < j_0 + N$ and $m_{i,j_0+N} \in 3\mathbb{Z}$.

Note that the set of all entries of the singularity diagram is the union of these subsets. Then we can see that the exceptional curve $C^{t,k'}$ obtained by the blow-up at $x^{i,j}$, $j_0 + 1 < j < j_0 + N$

does not contribute to $j''_{0,1}(F_p)$ from Lemma 1.14. Hence it contains at most $2a - 1$ points among $x_1, \dots, x_{j''_{0,1}(F_p)}$.

(ii) Assume that $C^{t,k}$ contributes to $j''_{0,a}(F_p)$. Then $C^{t,k}$ is blown up $3a - 2$ times when it is a (-2) -curve or $3a - 3$ times when it is a singular rational curve. Hence it contains at most $2a - 2$ points among $x_1, \dots, x_{j''_{0,1}(F_p)}$ by the same argument as in (i).

(iii) Assume that $C^{t,k}$ contributes to $j''_{1,a}(F_p)$. Then $C^{t,k}$ is blown up $3a$ times. Hence it contains at most $2a$ points among $x_1, \dots, x_{j''_{0,1}(F_p)}$ by the same argument as in (i).

We estimate $j''_{0,1}(F_p)$ from (i), (ii), (iii) as follows.

(a) If $D^t(p) = 0$ and $j''_{0,a}(F_p) = 0$ for any $a \geq 2$, then we have shown that $j''_{0,1}(F_p) \leq 2$ in Example 1.15 (2).

(b) If $D^t(p) = 0$ and $j''_{0,a}(F_p) > 0$ for some $a \geq 2$, then x_i is the point to which $C^{t,1}$ contracts or contained in some $C^{t,k}$ which contributes to $j''_{0,a}(F_p)$ for some $a \geq 2$. Hence we have

$$j''_{0,1}(F_p) \leq 1 + \sum_{a \geq 2} (2a - 1) j''_{0,a}(F_p).$$

(c) If $D^t(p) \neq 0$, then x_i is contained in some $C^{t,k}$ which does not contribute to $j''_{0,1}(F_p)$. Hence we have

$$j''_{0,1}(F_p) \leq \sum_{a \geq 2} (2a - 2) j''_{0,a}(F_p) + \sum_{a \geq 1} 2a j''_{1,a}(F_p) + \sum_{a \geq 2} (2a - 1) j''_{0,a}(F_p).$$

From (a), (b) and (c), we have

$$j''_{0,1}(F_p) \leq \bar{\eta}_p + \eta''_p + \sum_{a \geq 2} (2a - 2) j''_{0,a}(F_p) + \sum_{a \geq 1} 2a j''_{1,a}(F_p) + \sum_{a \geq 2} (2a - 1) j''_{0,a}(F_p)$$

by summing up for $t = 1, \dots, \eta_p$. Combining this with $\bar{\eta}_p \leq \eta''_p$, the claim (1) follows.

Suppose $n = 2$. Let $x^{t,k}$ be the point to which $C^{t,k}$ is contracted and $m^{t,k}$ the multiplicity of R at $x^{t,k}$. If $D^t(p) = 0$, then $x^{t,k}$, $k \geq 2$ is contained in $C^{t,k'}$ for some $k' < k$. Otherwise, $x^{t,1}$ is also contained in $C^{t,k'}$ for some k' . If $C^{t,k}$ is smooth, a singularity with odd multiplicity which is not contained in $C^{t,k'}$ for any $k' > k$ corresponds to an entry $(x_{i,j}, m_{i,j})$ of the singularity diagram $\mathcal{D}^{t,k}$ of $C^{t,k}$ satisfying that $m_{i,j-1}$ is even if $j > 1$, and $m_{i,j}$ is odd and then corresponds to a subset of entries of the diagram satisfying (*). For a curve $C^{t,k}$, we consider how many such subsets of entries of its singularity diagram there are.

(iv) If $C^{t,k}$ contributes to $j''_{0,a}(F_p)$, then $C^{t,k}$ is blown up $2a - 1$ times. Then the singularity diagram of $C^{t,k}$ has at most $a - 1$ subsets satisfying (*).

(v) If $C^{t,k}$ contributes to $j''_{0,a}(F_p)$ and it is a (-2) -curve, then $C^{t,k}$ is blown up $2a - 2$ times. Then the singularity diagram of $C^{t,k}$ has at most $a - 1$ subsets satisfying (*).

(vi) If $C^{t,k}$ contributes to $j''_{0,a}(F_p)$ and it is a singular rational curve, then $C^{t,k}$ is blown up $2a - 3$ times. Considering the singularity diagram of the proper transform of $C^{t,k}$ by the blow-up at its singular point, $C^{t,k}$ has at most $a - 1$ singularities with odd multiplicity which is not contained in $C^{t,k'}$ for any $k' > k$.

(vii) If $C^{t,k}$ contributes to $j_{1,a}''(F_p)$, then $C^{t,k}$ is blown up $2a$ times. Then the singularity diagram of $C^{t,k}$ has at most a subsets satisfying (*).

We estimate $j''(F_p)$ using (iv), (v), (vi) and (vii) as follows.

(d) If $D^t(p) = 0$, then the number of singularities with odd multiplicity appearing in $\{C^{t,k}\}_k$ is $j'''(F_p) - 1$. Hence we have

$$j'''(F_p) - 1 \leq \sum_{a \geq 2} (a-1)j_{0,a}'''(F_p).$$

(e) If $D^t(p) \neq 0$, then the number of singularities with odd multiplicity appearing in $\{C^{t,k}\}_k$ is $j''(F_p)$. Hence we have

$$j''(F_p) \leq j_{0,1,\text{odd}}''(F_p) + \sum_{a \geq 3} (a-1)j_{0,a}''(F_p) + \sum_{a \geq 1} a j_{1,a}''(F_p) + \sum_{a \geq 2} (a-1)j_{0,a}'''(F_p).$$

From (d) and (e), we have

$$j''(F_p) \leq \eta_p'' + j_{0,1,\text{odd}}'(F_p) + \sum_{a \geq 2} (a-1)j_{0,a}'(F_p) + \sum_{a \geq 1} a j_{1,a}'(F_p) + \sum_{a \geq 2} (a-1)j_{0,a}''(F_p)$$

by summing up for $t = 1, \dots, \eta_p$. Combining this with (1.12) and (1.14), the claim (2) follows. $\square$

Lemma 3.12. (1) If $n = 3$ and $j_{0,1}'(F_p) \neq 0$, then we have

$$\chi_\varphi(F_p) \geq \frac{1}{12}(j_{0,1}'(F_p) + 1).$$

(2) If $n = 2$, then the following hold.


(2, i) If Γ_p is a singular fiber not of type $({}_m\text{I}_k)$, (I_k^*) , (II^*) , (III^*) , (IV^*) , then we have


$$\chi_\varphi(F_p) \geq \frac{1}{12}(2j_{0,1}'(F_p) + j_{0,2}'(F_p) + j_{0,3}'(F_p) + 1).$$

Moreover, all the cases where Γ_p is a singular fiber of type $({}_m\text{I}_k)$, (II^*) , (III^*) , (IV^*) and

$$\chi_\varphi(F_p) < \frac{1}{12}(2j_{0,1}'(F_p) + j_{0,2}'(F_p) + j_{0,3}'(F_p) + 1)$$

are as follows.





where, in the dual graphs of Γ_p , the symbol $\circ$, $\bullet$, $\star$ respectively denotes a (-2) -curve not contained in R , contributing to $j'_{0,1}(F_p)$, contributing to $j'_{0,2}(F_p)$ or $j'_{0,3}(F_p)$. In these cases, we have

$$\chi_\varphi(F_p) = \frac{1}{12}(2j'_{0,1}(F_p) + j'_{0,2}(F_p) + j'_{0,3}(F_p) - 1)$$

when Γ_p is of type (III*) and $j'_{0,2}(F_p) = j'_{0,3}(F_p) = 0$ and


$$\chi_\varphi(F_p) = \frac{1}{12}(2j'_{0,1}(F_p) + j'_{0,2}(F_p) + j'_{0,3}(F_p))$$

otherwise.

(2, ii) If Γ_p is a singular fiber of type (I_k^*) , then

$$\chi_\varphi(F_p) \geq \frac{1}{12}(2j'_{0,1}(F_p) + j'_{0,2}(F_p) + j'_{0,3}(F_p) - 2)$$

with equality holding if and only if Γ_p and R satisfies the condition indicated in the following figure.



Proof. (1) Suppose that $n = 3$ and $j'_{0,1}(F_p) \neq 0$. If Γ_p is not of type (mI_k) , the claim is clear. Thus we may assume that Γ_p is of type (mI_k) . If $\chi_\varphi(F_p) = j'_{0,1}(F_p)/12$, then any component of Γ_p contributes to $j'_{0,1}(F_p)$ and contains at least 2 singular points of R , which is a contradiction.

(2) Suppose that $n = 2$. Any irreducible component C of Γ_p contributing to $j'_{0,1}(F_p)$ has no singular points of R . Thus any component of Γ_p intersecting with the curve C is not contained in R . From this observation and the classification of singular fibers of elliptic surfaces, the claims (2,i) and (2,ii) follow by an easy combinatorial argument. $\square$

Now, we are ready to prove Theorem 3.1.

Proof of Theorem 3.1. Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(g, 1, n)$. From Lemma 3.5, we have

$$\begin{aligned}
e_f(F_p) - \mu\chi_f(F_p) &= A_n\alpha_0^+(F_p) + \sum_{k \geq 1} (n - \mu'k) \alpha'_k(F_p) + n \sum_{k \geq 1} \alpha''_k(F_p) \\
&\quad - \mu' \sum_{k \geq 1} k\alpha''_k(F_p) - \sum_{a \geq 1} (2A_n + \delta_{a=1}) j_{0,a}(F_p) + C_n\chi_\varphi(F_p) - \frac{r\mu'}{n} \nu(F_p) \\
&= A_n\alpha_0^+(F_p) + \sum_{k \geq 1} (n - \mu'k) \alpha'_k(F_p) + C_n\chi_\varphi(F_p) \\
&\quad - \frac{r\mu'}{n} \left(1 - \frac{1}{m}\right) - \mu'\gamma_p + n\eta_p'' + \sum_{a \geq 1} (an^2 - a\mu') j'_{1,a}(F_p) \\
&\quad + \sum_{a \geq 1} (n(an - 2 - \delta_{mI_1, \mathbb{I}}) - a\mu' - 2A_n - \delta_{a=1}) j'_{0,a}(F_p) \\
&\quad + \sum_{a \geq 1} (n(an - 1) - a\mu' - 2A_n - \delta_{a=1}) j''_{0,a}(F_p) \\
&\quad - \mu' \sum_{k \geq 1} k\alpha_k^{n\mathbb{Z}+1}(F_p) - \sum_{k \geq 1} (n - \mu'k) \iota_k(F_p) - \sum_{k \geq 1} (n - \mu'k) \kappa_k(F_p),
\end{aligned}$$

where

$$A_n := n - 1 - \frac{(n-1)(2n-1)}{12n} \mu, \quad C_n := 12n - \left(\frac{r}{12n} (n-1)(n+1) + n \right) \mu$$

and $\mu' := (n-1)(n+1)\mu/12$. Combining Lemma 3.6 with the above equality, we have

$$\begin{aligned}
& e_f(F_p) - \mu \chi_f(F_p) \\
& \geq A_n \alpha_0^+(F_p) + \sum_{k \geq 1} (n - \mu' k) \alpha'_k(F_p) + C_n \chi_\varphi(F_p) - \frac{r\mu'}{n} \left(1 - \frac{1}{m}\right) \\
& \quad - \mu' \left(\frac{r}{n} - j'_{0,1}(F_p) \delta_{n=2} - \delta_{mI_1, \Pi} \right) \delta_{\eta'_p \neq 0} + (n - \mu') (\eta'_p - \delta_{\text{cyc}}) + \left(2n - \frac{r\mu'}{n} \right) \eta''_p \\
(3.5) \quad & + \sum_{a \geq 1} (n(an - 3 - \delta_{mI_1, \Pi}) - (a - 1)\mu' - 2A_n - \delta_{a=1}) j'_{0,a}(F_p) \\
& + \sum_{a \geq 1} (n(an - 2) - (a - 1)\mu' - 2A_n - \delta_{a=1}) j''_{0,a}(F_p) \\
& + \sum_{a \geq 1} (n(an - 1) - (a - 1)\mu') j'_{1,a}(F_p) - \mu' \sum_{k \geq 1} k \alpha_k^{n\mathbb{Z}+1}(F_p) - \sum_{k \geq 1} (n - \mu' k) \kappa_k(F_p).
\end{aligned}$$

Assume that $A_n \geq 0$ and $C_n \geq 0$. We obtain by using Lemmas 3.4 and 3.8 (1) that

$$\begin{aligned}
& e_f(F_p) - \mu \chi_f(F_p) \\
& \geq \sum_{k \geq 1} (n - \mu' k) \alpha'_k(F_p) + \frac{r(m-1)(n-1)(4-\mu)}{4m} + C_n \chi_\varphi(F_p) - \mu' \left(\frac{r}{n} - \delta_{mI_1, \Pi} \right) \delta_{\eta'_p \neq 0} \\
& \quad + A_n \beta_p + (n - \mu' - (n-2)A_n) (\eta'_p - \delta_{\text{cyc}}) + \left(2n - \frac{r\mu'}{n} - (n-2)A_n \right) \eta''_p \\
(3.6) \quad & + \sum_{a \geq 1} ((n-4)A_n + n(an - 3 - \delta_{mI_1, \Pi}) - (a-1)\mu' - (1 - \delta_{n=2}\mu')\delta_{a=1}) j'_{0,a}(F_p) \\
& + \sum_{a \geq 1} \left((n-4)A_n + n(an - 2) - \mu' \left(a + \frac{r}{n} - 2 \right) - \delta_{a=1} \right) j''_{0,a}(F_p) \\
& + \sum_{a \geq 1} ((n-2)A_n + n(an - 1) - (a-1)\mu') j'_{1,a}(F_p) \\
& + \left(2(n-2)A_n - n + \left(\frac{r}{n} - 1 \right) \mu' \right) \kappa(F_p) - \mu' \left(\frac{r}{n} - 2 \right) \kappa^{(3)}(F_p) - \mu' \alpha_{\frac{r}{nm}}^{n\mathbb{Z}+1}(F_p).
\end{aligned}$$

We put

$$\mu = \begin{cases} \frac{12n^2}{r(n-1)(n+1)}, & \text{if } n \geq 4, \\ \frac{24}{4r-13}, & \text{if } n = 3, \\ \frac{4}{r-2}, & \text{if } n = 2. \end{cases}$$

(i) We assume $n \geq 4$. We write $r = kn$. The coefficient of η'_p is

$$n - \frac{1}{12}(n-1)(n+1)\mu - (n-2)A_n = -(n^2 - 4n + 2) + \frac{n^2 - 6n + 2}{k(n+1)} < 0.$$

The coefficient of η''_p is

$$2n - \frac{(n-1)(n+1)r\mu}{12n} - (n-2)A_n = -(n^2 - 4n + 2) + \frac{(n-2)(2n-1)}{k(n+1)}.$$

It is negative if $n \geq 5$ or $k \geq 2$. Note that $\eta''_p = 0$ if $k < n - 1$ since the multiplicity m' of a singular point of type $n\mathbb{Z} + 1$ satisfies $(n-1)^2 \leq m' \leq r - n + 1$. Thus, we may not consider the case where $n = 4$ and $k = 1$. Using $\eta'_p \leq j'(F_p)$ and $\eta''_p \leq j''(F_p)$, (3.6) is greater than or equal to

$$\begin{aligned} & \sum_{k \geq 1} (n - \mu'k) \alpha'_k(F_p) + \frac{r(m-1)(n-1)(4-\mu)}{4m} + C_n \chi_\varphi(F_p) + A_n \beta_p \\ & - \mu' \left(\frac{r}{n} - \delta_{mI_1, \mathbb{I}} \right) \delta_{\eta'_p \neq 0} + ((n-2)A_n - n + \mu') \delta_{\text{cyc}} + \sum_{a \geq 1} (an^2 - a\mu') j'_{1,a}(F_p) \\ (3.7) \quad & + \sum_{a \geq 1} (n(an - 2 - \delta_{mI_1, \mathbb{I}}) - a\mu' - 2A_n - \delta_{a=1}) j'_{0,a}(F_p) \\ & + \sum_{a \geq 1} \left(an^2 - \mu' \left(a + \frac{2r}{n} - 2 \right) - 2A_n - \delta_{a=1} \right) j''_{0,a}(F_p) \\ & + \left(2(n-2)A_n - n + \left(\frac{r}{n} - 1 \right) \mu' \right) \kappa(F_p) - \mu' \left(\frac{r}{n} - 2 \right) \kappa^{(3)}(F_p). \end{aligned}$$

Since $2(n-2)A_n - n + \mu' > 0$, we have

$$\begin{aligned} & \left(2(n-2)A_n - n + \left(\frac{r}{n} - 1 \right) \mu' \right) \kappa(F_p) - \mu' \left(\frac{r}{n} - 2 \right) \kappa^{(3)}(F_p) \\ & \geq (2(n-2)A_n - n + \mu') \kappa(F_p) \\ & \geq 0. \end{aligned}$$

The coefficient of $j'_{0,1}(F_p)$ in (3.7) is

$$\begin{aligned} & n(n-2 - \delta_{mI_1, \mathbb{I}}) - \mu' - 2A_n - \delta_{a=1} \\ (3.8) \quad & = n^2 - (4 + \delta_{mI_1, \mathbb{I}})n + 1 - \frac{(n-1)(n-2)}{k(n+1)}. \end{aligned}$$

If $\delta_{mI_1, \mathbb{I}} = 0$, (3.8) is negative if and only if $n = 4$ and $k = 1$. If $\delta_{mI_1, \mathbb{I}} = 1$ and $n \geq 5$, (3.8) is non-negative. If $\delta_{mI_1, \mathbb{I}} = 1$ and $n = 4$, (3.8) is $-3 - 6/5k < 0$. Note that $j'(F_p) = 0$ if $k = 1$ since $r = n$ and any singularity of R has the multiplicity n . Thus, we may not consider the case where $\delta_{mI_1, \mathbb{I}} = 0$, $n = 4$ and $k = 1$. We can check that the coefficient of $j'_{0,a}(F_p)$ in (3.7) is positive for $a \geq 2$. Moreover, we also can check that the coefficient of $j''_{0,a}(F_p)$ in (3.7) is positive for $a \geq 1$.

(i,1) We assume that $\eta'_p = 0$. Then, clearly (3.7) is non-negative.

(i,2) We assume that $\eta'_p \neq 0$, $\chi_\varphi(F_p) \geq 1/6$ (i.e. Γ_p is a singular fiber not of type (mI_1)) and $\iota^{(3)}(F_p) = 0$. Then we have

$$C_n \chi_\varphi(F_p) \geq \frac{1}{6} C_n = \frac{n}{6} \left(11 - \frac{12n}{k(n-1)(n+1)} \right).$$

The coefficient of $\delta_{\eta'_p \neq 0}$ is

$$\mu' \left(\frac{r}{n} - \delta_{\text{mI}_1, \text{II}} \right) = -n + \frac{n}{k} \delta_{\text{mI}_1, \text{II}}.$$

If $n \neq 4$ or $\delta_{\text{mI}_1, \text{II}} = 0$ or $j'_{0,1}(F_p) = 0$, then (3.7) is positive since

$$\frac{n}{6} \left(11 - \frac{12n}{k(n-1)(n+1)} \right) - n = \frac{n}{6} \left(5 - \frac{12n}{k(n-1)(n+1)} \right) > 0.$$

If $n = 4$ and $\delta_{\text{mI}_1, \text{II}} = 1$ and $j'_{0,1}(F_p) \neq 0$ (we denote this condition by $(\#)$), then we have $j'_{0,1}(F_p) = 1$ and then (3.7) is positive since

$$\frac{n}{6} \left(11 - \frac{12n}{k(n-1)(n+1)} \right) - n + \frac{n}{k} - 3 - \frac{6}{5k} = \frac{1}{3} + \frac{26}{15k} > 0.$$

(i,3) We assume that $\eta'_p \neq 0$ and $\chi_\varphi(F_p) = 1/12$ (i.e. Γ_p is a singular fiber of type (mI_1)). Let m_1 be the multiplicity of the singular point x_1 of R which is singular for Γ_p . If $m_1 \in n\mathbb{Z}$, then x_1 contributes $m_1 - 2$ to $\alpha_0^+(F_p)$ (Note that x_1 is a 1-vertical type singularity). In particular, x_1 contributes at least $n - 2$ to $\alpha_0^+(F_p)$. If the condition $(\#)$ does not hold, (3.7) is positive since

$$\frac{n}{12} \left(11 - \frac{12n}{k(n-1)(n+1)} \right) - n + (n-2)A_n = n^2 - \frac{37}{12}n + 2 - \frac{n^2 + (n-2)(n-1)(2n-1)}{k(n-1)(n+1)}$$

increases monotonically with respect to n and

$$\frac{17}{3} - \frac{58}{15k} > 0$$

when $n = 4$. If the condition $(\#)$ holds, (3.7) is also positive since

$$\frac{17}{3} - \frac{58}{15k} + \frac{4}{k} - 3 - \frac{6}{5k} = \frac{8}{3} - \frac{16}{15k} > 0.$$

(i,4) We assume that $\eta'_p \neq 0$ and $\chi_\varphi(F_p) = 0$ (i.e. Γ_p is a smooth elliptic curve). Then (3.7) is positive since $j'_{1,\bullet}(F_p) = 1$.

(i,5) We assume that $\iota^{(3)}(F_p) = 1$. From Lemma 3.3, we may consider the following 3 cases.

(i,5,II) If Γ_p is a singular fiber of type (II) , then $\beta_p = n - 7$, $\chi_\varphi(F_p) = 1/6$ and $\kappa(F_p) = \kappa^{(2)}(F_p) \geq 1$. From the argument in (i,2), it is sufficient to show that

$$A_n \beta_p + \left(2(n-2)A_n - n + \left(\frac{r}{n} - 1 \right) \mu' \right) \kappa(F_p) > 0.$$

This inequality is true, since

$$\begin{aligned}
& (n-7)A_n + 2(n-2)A_n - n + \left(\frac{r}{n} - 1\right)\mu' \\
&= (3n-11) \left(n-1 - \frac{2n-1}{k(n+1)}\right) - \frac{n}{k} \\
&> 0.
\end{aligned}$$

Note that $k \geq n-1$ since $j''(F_p) \neq 0$.

(i,5,III) If Γ_p is a singular fiber of type (III), then $\beta_p = -n-1$, $\chi_\varphi(F_p) = 1/4$ and $\kappa(F_p) = \kappa^{(2)}(F_p) \geq 1$. Since $j''(F_p) \geq 1$ and the coefficient of $j''_{0,a}(F_p)$ is greater than 1 for any $a \geq 1$, it is sufficient to show that

$$1 + A_n\beta_p + \frac{1}{12}C_n + \left(2(n-2)A_n - n + \left(\frac{r}{n} - 1\right)\mu'\right)\kappa(F_p) \geq 0.$$

The left hand side of it is greater than or equal to

$$\begin{aligned}
& 1 - (n+1)A_n + \frac{1}{12}C_n + 2(n-2)A_n - n + \left(\frac{r}{n} - 1\right)\mu' \\
(3.9) \quad &= n^2 - \frac{61}{12}n + 6 - \frac{3n^3 - 12n^2 + 15n - 5}{k(n-1)(n+1)}
\end{aligned}$$

and (3.9) increases monotonically with respect to n . If $n = 4$, (3.9) is $5/3 - 11/3k > 0$. Note that $k \geq n-1$ since $j''(F_p) \neq 0$.

(i,5,IV) If Γ_p is a singular fiber of type (IV), then $\beta_p = -2$, $\chi_\varphi(F_p) = 1/3$ and $j'_{0,\bullet}(F_p) \geq 3$. Thus, it is sufficient to show that $A_n\beta_p + C_n/6 + 3 \cdot (3.8)$ is positive. By a computation, this is equal to

$$3n^2 - \frac{73}{6}n + 5 - \frac{3n^3 - 14n^2 + 21n - 8}{k(n-1)(n+1)}$$

and we can check that it is positive.

From (i,1) through (i,5), we have $e_f(F_p) - \mu\chi_f(F_p) \geq 0$ for $n \geq 4$. On the other hand, if $n = 3$ and $g = 4$, one can easily classify all singular fibers of primitive cyclic covering fibrations of type $(4, 1, 3)$ because R has no singularities of multiplicity greater than 3, and check $e_f(F_p) \geq (9/2)\chi_f(F_p)$ for any fiber germ F_p . Thus, Theorem 3.1 (1) follows.

(ii) We assume $n = 3$ and $g > 4$. The coefficient of $j''_{0,1}(F_p)$ in (3.6) is

$$-\left(\frac{2}{9}r - \frac{17}{18}\right)\mu < 0.$$

Applying Lemma 3.11 (1) to the term of $j''_{0,1}(F_p)$, (3.6) is greater than or equal to

$$\begin{aligned}
(3.10) \quad & \sum_{k \geq 1} \left(3 - \frac{2}{3} \mu k \right) \alpha'_k(F_p) + \frac{r(m-1)(4-\mu)}{2m} + \left(36 - \left(\frac{2}{9} r + 3 \right) \mu \right) \chi_\varphi(F_p) \\
& + \left(2 - \frac{5}{18} \mu \right) \beta_p - \frac{2}{3} \mu \left(\frac{r}{3} - \delta_{mI_1, \mathbb{II}} \right) \delta_{\eta'_p \neq 0} + \left(1 - \frac{7}{18} \mu \right) (\eta'_p - \delta_{\text{cyc}}) \\
& + \sum_{a \geq 2} \left(9a - 11 - 3\delta_{mI_1, \mathbb{II}} - \left(\frac{4}{9}(a-1)r - \frac{11}{9}a + \frac{17}{18} \right) \mu \right) j'_{0,a}(F_p) \\
& + \left(-3 + \frac{5}{18} \mu \right) j'_{0,1}(F_p) + \sum_{a \geq 2} \left(9a - 8 - \left(\frac{4}{9}ar - \frac{11}{9}a - \frac{2}{3} \right) \mu \right) j''_{0,a}(F_p) \\
& + \sum_{a \geq 1} \left(9a - 1 - \left(\frac{4}{9}ar - \frac{11}{9}a - \frac{7}{18} \right) \mu \right) j'_{1,a}(F_p) \\
& + \left(1 + \left(\frac{2}{9}r - \frac{11}{9} \right) \mu \right) \kappa(F_p) - \frac{2}{3} \mu \left(\frac{r}{3} - 2 \right) \kappa^{(3)}(F_p).
\end{aligned}$$

We remark that the term of η''_p vanishes by the definition of μ and

$$\begin{aligned}
& \left(1 + \left(\frac{2}{9}r - \frac{11}{9} \right) \mu \right) \kappa(F_p) - \frac{2}{3} \mu \left(\frac{r}{3} - 2 \right) \kappa^{(3)}(F_p) \\
& \geq \left(1 + \frac{1}{9} \mu \right) \kappa(F_p) \\
& \geq 0.
\end{aligned}$$

We can check that the coefficient of $j'_{0,a}(F_p)$ (resp. $j''_{0,a}(F_p)$, $j'_{1,a}(F_p)$) in (3.10) are positive for $a \geq 2$ (resp. $a \geq 2$, $a \geq 1$).

(ii,1) Assume that $\eta'_p = 0$. Then (3.10) is non-negative.

(ii,2) Assume that $\eta'_p \neq 0$ and $j_{0,1}^{tt}(F_p) = 4$ for some t . From Lemma 3.10 (1,iii), it follows that Γ_p is a singular fiber of type (I_k^*) for some k , $\chi_\varphi(F_p) = (k+6)/12$, $r \geq 9$, $\Gamma_p \subset R$, $\eta'_p = 1$, $j'_{0,1}(F_p) = 4$ and $\delta_{\text{cyc}} = 0$. Considering the terms of $\chi_\varphi(F_p)$, $\delta_{\eta'_p \neq 0}$, η'_p and $j'_{0,1}(F_p)$, (3.10) is greater than or equal to

$$\begin{aligned}
& \left(36 - \left(\frac{2}{9}r + 3 \right) \mu \right) \frac{k+6}{12} - \frac{2}{9}r\mu + \left(1 - \frac{7}{18} \mu \right) + 4 \left(-3 + \frac{5}{18} \mu \right) \\
& = 7 - \frac{8(3r+2)}{3(4r-13)}.
\end{aligned}$$

This is positive since $r \geq 9$.

(ii,3) Assume that $\eta'_p \neq 0$, $\iota^{(3)}(F_p) = 0$, $r \geq 9$ and $j_{0,1}^{tt}(F_p) \geq 3$ for any t . Then

$$\frac{1}{3} j'_{0,1}(F_p) \leq \eta'_p - \delta_{\text{cyc}}$$

from Lemma 3.10 (1,i), (1,ii). From this and Lemma 3.12 (1), (3.10) is greater than or equal to

$$\left(36 - \left(\frac{2}{9}r + 3\right)\mu\right) \frac{j'_{0,1}(F_p) + 1}{12} - \frac{2}{9}r\mu + \frac{1}{3} \left(1 - \frac{7}{18}\mu\right) j'_{0,1}(F_p) + \left(-3 + \frac{5}{18}\mu\right) j'_{0,1}(F_p).$$

One can check by a computation that this is positive since $r \geq 9$.

(ii,4) Assume that $\eta'_p \neq 0$, $\iota^{(3)}(F_p) = 0$, $r = 6$ and $\chi_\varphi(F_p) \geq (j'_{0,1}(F_p) + 2)/12$. By the same argument as in (ii,3), (3.10) is greater than or equal to

$$\begin{aligned} & \left(36 - \left(\frac{2}{9}r + 3\right)\mu\right) \frac{j'_{0,1}(F_p) + 2}{12} - \frac{2}{9}r\mu + \frac{1}{3} \left(1 - \frac{7}{18}\mu\right) j'_{0,1}(F_p) + \left(-3 + \frac{5}{18}\mu\right) j'_{0,1}(F_p) \\ &= -\frac{13}{99}j'_{0,1}(F_p) + \frac{50}{33}. \end{aligned}$$

On the other hand, one sees that $j'_{0,1}(F_p) \leq 6$ since $r = 6$. Then

$$-\frac{13}{99}j'_{0,1}(F_p) + \frac{50}{33} > 0.$$

(ii,5) Assume that $\eta'_p \neq 0$, $\iota^{(3)}(F_p) = 0$, $r = 6$ and $\chi_\varphi(F_p) < (j'_{0,1}(F_p) + 2)/12$. If $j'_{0,1}(F_p) = 0$, then (3.10) is positive since $j'(F_p) \neq 0$. Then there are the following two cases only.

(ii,5,I₂) Γ_p is a singular fiber of type ($_m\text{I}_2$) and only one component of Γ_p is contained in R and brown up just once.

(ii,5,I₃) Γ_p is a singular fiber of type ($_m\text{I}_3$) and only two component of Γ_p are contained in R and brown up just once at the intersection point of these.

In both cases, we can see that

$$\gamma_p \leq \left(\frac{r}{n} - 1\right) \delta_{\eta'_p \neq 0} + \left(\frac{r}{n} - 1\right) \eta''_p$$

from the proof of Lemma 3.6, since the component of Γ_p not contained in R intersects with R_h . Thus, it is sufficient to show that

$$\left(36 - \left(\frac{2}{9}r + 3\right)\mu\right) \frac{j'_{0,1}(F_p) + 1}{12} - \frac{2}{3}\mu \left(\frac{r}{3} - 1\right) + \left(1 - \frac{7}{18}\mu\right) + \left(-3 + \frac{5}{18}\mu\right) j'_{0,1}(F_p)$$

is positive. This is equal to $(-2j'_{0,1}(F_p) + 10)/11 > 0$ by a computation.

(ii,6) Assume that $\iota^{(3)}(F_p) = 1$. From Lemma 3.3, we may consider the following 3 cases.

(ii,6,II) If Γ_p is a singular fiber of type (II), then we have $\chi_\varphi(F_p) = 1/6$, $\beta_p = -4$, $\delta_{\text{cyc}} = 0$, $j'_{0,1}(F_p) = 0$, $j'_{0,\bullet}(F_p) = 1$, $j''_{0,\bullet}(F_p) \geq 2$ and $\kappa(F_p) = \kappa^{(2)}(F_p) \geq 1$. One can see easily that (3.10) is positive by a computation.

(ii,6,III) If Γ_p is a singular fiber of type (III), then we have $\chi_\varphi(F_p) = 1/4$, $\beta_p = -4$, $\delta_{\text{cyc}} = 0$, $j'_{0,1}(F_p) = 0$, $j'_{0,\bullet}(F_p) = 2$, $j''_{0,\bullet}(F_p) \geq 1$ and $\kappa(F_p) = \kappa^{(2)}(F_p) \geq 1$. One can see easily that (3.10) is positive by a computation.

(ii,6,IV) If Γ_p is a singular fiber of type (IV), then we have $\chi_\varphi(F_p) = 1/3$, $\beta_p = -2$, $\delta_{\text{cyc}} = 0$, $j'_{0,\bullet}(F_p) = 3$ and $\kappa^{(3)}(F_p) = 0$. If $j'_{0,1}(F_p) \leq 2$, then we can check that (3.10) is positive.

Suppose that $j'_{0,1}(F_p) = 3$. Then any component of Γ_p is brown up only once. Thus, the multiplicity of the singularity of R on Γ_p is $r/3 + 3 \geq 6$. In particular, $r \geq 9$. Since this singularity is a 3-vertical $3\mathbb{Z}$ type singularity, we have

$$\alpha_0^+(F_p) \geq \iota(F_p) + 2\kappa(F_p) + \beta_p + 3$$

from the proof of Lemma 3.4. Then it suffices to show that

$$\left(2 - \frac{5}{18}\mu\right) + \left(36 - \left(\frac{2}{9}r + 3\right)\mu\right)\frac{1}{3} - \frac{2}{9}r\mu + \left(1 - \frac{7}{18}\mu\right) + 3\left(-3 + \frac{5}{18}\mu\right)$$

is positive. This is equal to

$$6 - \frac{4(16r + 45)}{9(4r - 13)} > 0.$$

From (ii,1) through (ii,6), we have $e_f(F_p) - \mu\chi_f(F_p) \geq 0$. Thus, Theorem 3.1 (2) follows.

(iii) We assume $n = 2$ and $g \geq 3$. From Lemmas 3.4 and 3.8 (2), (3.5) is greater than or equal to

$$\begin{aligned} & \sum_{k \geq 1} \left(2 - \frac{1}{4}\mu k\right) \alpha'_k(F_p) + \frac{r(m-1)(4-\mu)}{4m} + \left(24 - \left(\frac{r}{8} + 2\right)\mu\right) \chi_\varphi(F_p) \\ & - \frac{1}{4}\mu \left(\frac{r}{2} - j'_{0,1}(F_p) - \delta_{mI_1, \mathbb{I}}\right) \delta_{\eta'_p \neq 0} + \left(2 - \frac{1}{4}\mu\right) (\eta'_p - \delta_{\text{cyc}}) + \left(4 - \frac{1}{8}r\mu\right) \widehat{\eta}_p \\ & + \sum_{a \geq 1} \left(4a - 8 - 2\delta_{mI_1, \mathbb{I}} - \delta_{a=1} - \frac{1}{4}(a-2)\mu\right) j'_{0,a}(F_p) \\ (3.11) \quad & + \sum_{a \geq 2} \left(4a - 6 - \frac{1}{4}\left(a + \frac{r}{2} - 3\right)\mu\right) j''_{0,a}(F_p) + \sum_{a \geq 1} \left(4a - 2 - \frac{1}{4}(a-1)\mu\right) j'_{1,a}(F_p) \\ & + \sum_{k \geq 1} \left(1 - \frac{1}{8}(r-2)\mu - \frac{1}{4}\mu k\right) \alpha^{\text{tr}}_{(2k+1 \rightarrow 2k+1)}(F_p) - \frac{1}{4}\mu \alpha^{\frac{2\mathbb{Z}+1}{2m}}_r(F_p) \\ & + \sum_{k \geq 1} \left(1 - \frac{1}{8}(r-2)\mu + \left(2 - \frac{1}{2}\mu\right)k\right) \alpha^{\text{co},0}_{(2k+1 \rightarrow 2k+1)}(F_p) - \left(2 - \frac{1}{8}(r-2)\mu\right) \kappa(F_p) \\ & + \sum_{k \geq 1} \left(-3 + \frac{1}{4}\mu + \left(2 - \frac{1}{2}\mu\right)k\right) \alpha^{\text{co},1}_{(2k+1 \rightarrow 2k+1)}(F_p) - \frac{1}{8}(r-4)\mu \kappa^{(3)}(F_p). \end{aligned}$$

The coefficients of $\alpha^{\text{tr}}_{(2k+1 \rightarrow 2k+1)}(F_p)$ and $\alpha^{\text{co},0}_{(2k+1 \rightarrow 2k+1)}(F_p)$ in (3.11) are non-negative and that of $\alpha^{\text{tr}}_{(r-1 \rightarrow r-1)}(F_p)$ is 0 by the definition of μ . The coefficient of $\alpha^{\text{co},1}_{(2k+1 \rightarrow 2k+1)}(F_p)$ in (3.11) is positive except for $k = 1$ and that of $\alpha^{\text{co},1}_{(3 \rightarrow 3)}(F_p)$ is $-1 - \mu/4 < 0$. The coefficient of $\kappa(F_p)$ in (3.11) is $-3/2$. Applying Lemmas 3.9 and 3.11 (2) to $(3/2)\kappa(F_p)$ and $(1 + \mu/4) \sum_{k \geq 1} \alpha^{\text{co},1}_{(2k+1 \rightarrow 2k+1)}(F_p)$, we see that (3.11) is greater than or equal to

$$\begin{aligned}
& \sum_{k \geq 1} \left(2 - \frac{1}{4} \mu k \right) \alpha'_k(F_p) + \frac{r(m-1)(4-\mu)}{4m} + \left(24 - \left(\frac{r}{8} + 2 \right) \mu \right) \chi_\varphi(F_p) \\
& - \frac{1}{4} \mu \left(\frac{r}{2} - \delta_{mI_1, \Pi} \right) \delta_{\eta'_p \neq 0} + \left(2 - \frac{1}{4} \mu \right) (\eta'_p - \delta_{\text{cyc}}) + \left(3 - \frac{1}{8} (r+2) \mu \right) \widehat{\eta}_p \\
& - \left(5 - \frac{1}{2} \mu \right) j'_{0,1}(F_p) - (1 + 2\delta_{mI_1, \Pi}) j'_{0,2,\text{even}}(F_p) - \left(2 + \frac{1}{4} \mu \right) j'_{0,2,\text{odd}}(F_p) \\
& - \left(2\delta_{mI_1, \Pi} + \frac{3}{4} \mu \right) j'_{0,3}(F_p) + \sum_{a \geq 4} \left(2a - 6 - 2\delta_{mI_1, \Pi} - \frac{1}{4} (2a-3) \mu \right) j'_{0,a}(F_p) \\
(3.12) \quad & + \sum_{a \geq 2} \left(2a - 3 - \frac{1}{4} \left(2a + \frac{r}{2} - 5 \right) \mu \right) j''_{0,a}(F_p) + \sum_{a \geq 1} \left(2a - \frac{1}{2} (a-1) \mu \right) j'_{1,a}(F_p) \\
& + \sum_{k \geq 1} \left(1 - \frac{1}{8} (r-2) \mu - \frac{1}{4} \mu k \right) \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{tr}}(F_p) + \left(1 - \frac{1}{4} \mu \right) \alpha_{\frac{r}{2m}}^{2\mathbb{Z}+1}(F_p) \\
& + \sum_{k \geq 1} \left(1 - \frac{1}{8} (r-2) \mu + \left(2 - \frac{1}{2} \mu \right) k \right) \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},0}(F_p) \\
& + \sum_{k \geq 1} \left(2 - \frac{1}{2} \mu \right) (k-1) \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},1}(F_p) - \frac{1}{8} (r-4) \mu \kappa^{(3)}(F_p),
\end{aligned}$$

where $j'_{0,2,\text{even}}(F_p) = j'_{0,2}(F_p) - j'_{0,2,\text{odd}}(F_p)$.

(iii,1) Assume that $\eta'_p = 0$. Then (3.12) is clearly non-negative.

(iii,2) Assume that $\eta'_p \neq 0$ and $j_{0,2,\text{odd}}^t(F_p) = 4$ for some t . From Lemma 3.10 (2,iii), Γ_p is a singular fiber of type (I_k^*) , $\Gamma_p \subset R$, $\kappa^{(3)}(F_p) = 0$, $\eta'_p = 1$, $\delta_{\text{cyc}} = 0$, $j'_{0,1}(F_p) = 0$ and $j'_{0,2,\text{odd}}(F_p) = 4$. Clearly we have $\chi_\varphi(F_p) = (j'_{0,2}(F_p) + j'_{0,3}(F_p) + 1)/12$. Considering the terms of $\chi_\varphi(F_p)$, $\delta_{\eta'_p \neq 0}$, η'_p , $j'_{0,2,\text{even}}(F_p)$, $j'_{0,2,\text{odd}}(F_p)$ and $j'_{0,3}(F_p)$, (3.12) is greater than or equal to

$$\begin{aligned}
& \left(24 - \left(\frac{r}{8} + 2 \right) \mu \right) \frac{1}{12} (j'_{0,2,\text{even}}(F_p) + j'_{0,3}(F_p) + 5) - \frac{1}{8} r \mu + \left(2 - \frac{1}{4} \mu \right) \\
& - j'_{0,2,\text{even}}(F_p) - 4 \left(2 + \frac{1}{4} \mu \right) - \frac{3}{4} \mu j'_{0,3}(F_p) \\
& = \left(1 - \left(\frac{r}{96} + \frac{1}{6} \right) \mu \right) j'_{0,2,\text{even}}(F_p) + \left(2 - \left(\frac{r}{96} + \frac{11}{12} \right) \mu \right) j'_{0,3}(F_p) + 4 - \left(\frac{17}{96} r + \frac{25}{12} \right) \mu,
\end{aligned}$$

which is positive.

(iii,3) Assume that $\eta'_p \neq 0$, $\kappa^{(3)}(F_p) = 0$, $\delta_{mI_1, \Pi} = 0$, $j_{0,2,\text{odd}}^t(F_p) \leq 3$ for any t and

$$\chi_\varphi(F_p) \geq \frac{1}{12} (2j'_{0,1}(F_p) + j'_{0,2}(F_p) + j'_{0,3}(F_p) + 1).$$

From Lemma 3.10 (2,i), (2,ii), we have $j'_{0,1}(F_p) + j'_{0,2,\text{odd}}(F_p)/3 \leq \eta'_p - \delta_{\text{cyc}}$. Then (3.12) is greater than or equal to

$$\begin{aligned}
& \left(24 - \left(\frac{r}{8} + 2\right)\mu\right) \frac{1}{12}(2j'_{0,1}(F_p) + j'_{0,2}(F_p) + j'_{0,3}(F_p) + 1) - \frac{1}{8}r\mu \\
& + \left(2 - \frac{1}{4}\mu\right) \left(j'_{0,1}(F_p) + \frac{1}{3}j'_{0,2,\text{odd}}(F_p)\right) - \left(5 - \frac{1}{2}\mu\right) j'_{0,1}(F_p) - j'_{0,2,\text{even}}(F_p) \\
& - \left(2 + \frac{1}{4}\mu\right) j'_{0,2,\text{odd}}(F_p) - \frac{3}{4}\mu j'_{0,3}(F_p) \\
& = \left(1 - \left(\frac{r}{48} + \frac{1}{12}\right)\mu\right) j'_{0,1}(F_p) + \left(\frac{2}{3} - \left(\frac{r}{96} + \frac{1}{2}\right)\mu\right) j'_{0,2,\text{odd}}(F_p) \\
& + \left(1 - \left(\frac{r}{96} + \frac{1}{6}\right)\mu\right) j'_{0,2,\text{even}}(F_p) + \left(2 - \left(\frac{r}{96} + \frac{11}{12}\right)\mu\right) j'_{0,3}(F_p) \\
& + 2 - \left(\frac{13}{96}r + \frac{1}{6}\right)\mu.
\end{aligned}$$

The coefficients of $j'_{0,1}(F_p)$, $j'_{0,2,\text{even}}(F_p)$, $j'_{0,3}(F_p)$ and the constant term are positive since $r \geq 4$, and $j_{0,2,\text{odd}}(F_p)$ is also positive for $r \geq 6$. Thus the above equation is positive when $r \geq 6$. If $r = 4$, then one can check by an easy computation that (3.12) is also positive, since $j'_{0,2,\text{odd}}(F_p) \leq 2$.

(iii,4) Assume that $\eta'_p \neq 0$, $\kappa^{(3)}(F_p) = 0$, $\delta_{m\text{I}_1, \text{II}} = 0$, $j_{0,2,\text{odd}}^{tt}(F_p) \leq 3$ for any t and

$$\chi_\varphi(F_p) < \frac{1}{12}(2j'_{0,1}(F_p) + j'_{0,2}(F_p) + j'_{0,3}(F_p) + 1).$$

From Lemma 3.12 (2), Γ_p is of type $({}_m\text{I}_k)$, (I_k^*) , (II^*) , (III^*) or (IV^*) . Considering the numbers $j'_{0,1}(F_p)$, $\chi_\phi(F_p)$ and Lemma 3.12 (2), we can see that (3.12) is positive by the same argument as in (iii,3) except Γ_p is of type (I_0^*) and $j'_{0,1}(F_p) = 4$.

Suppose that Γ_p is of type (I_0^*) and $j'_{0,1}(F_p) = 4$. Then the component not contributing to $j'_{0,1}(F_p)$ is a double component in Γ_p and intersects with R_h . Thus we have

$$\alpha_0^+(F_p) \geq \frac{r}{2} + \sum_{k \geq 1} 2k\alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co}}(F_p)$$

from the proof of Lemma 3.4. Then one can see that $e_f(F_p) - \mu\chi_\varphi(F_p)$ is positive by a computation.

(iii,5) Assume that $\eta'_p \neq 0$, $\kappa^{(3)}(F_p) = 0$ and $\delta_{m\text{I}_1, \text{II}} = 1$. Then $j'_{0,1}(F_p) = 0$, $j'(F_p) = 1$, $\eta'_p = 1$ and $\chi_\varphi(F_p) = 1/12$ or $1/6$. If $j'_{0,2,\text{even}}(F_p) = 1$, then $(\Gamma_p)_{\text{red}}$ is blown up just once. Thus, the multiplicity of the singularity of R which is singular for $(\Gamma_p)_{\text{red}}$ is even. Hence $\delta_{\text{cyc}} = 0$. Then we can check by a computation that (3.12) is positive. If $j'_{0,2,\text{even}}(F_p) = 0$, then we can also check by a computation that (3.12) is positive.

(iii,6) Assume that $\kappa^{(3)}(F_p) = 1$. From Lemma 3.3, we may consider the following 3 cases.

(iii,6,II) If Γ_p is a singular fiber of type (II), then $\eta'_p = 1$, $j'(F_p) = 1$, $j'_{0,a}(F_p) = 0$ for $a \leq 3$, $j''(F_p) - j''_{0,1}(F_p) \geq 3$ and $\chi_\varphi(F_p) = 1/6$. Considering the terms of $\delta_{\eta'_p \neq 0}$, η'_p , $\kappa^{(3)}(F_p)$, $j'_{0,a}(F_p)$, $j''_{0,a}(F_p)$ and $\chi_\varphi(F_p)$ in (3.12), we can check that (3.12) is positive.


(iii,6,III) If Γ_p is a singular fiber of type (III), then $\eta'_p = 1$, $j'(F_p) = 2$, $j'_{0,a}(F_p) = 0$ for $a \leq 2$, $j''(F_p) - j''_{0,1}(F_p) \geq 2$ and $\chi_\varphi(F_p) = 1/4$. Then we can also check that (3.12) is positive.

(iii,6,IV) If Γ_p is a singular fiber of type (IV), then $\eta'_p = 1$, $j'(F_p) = 3$, $j'_{0,1}(F_p) = j'_{0,2,\text{even}}(F_p) = 0$, $j''(F_p) - j''_{0,1}(F_p) \geq 1$ and $\chi_\varphi(F_p) = 1/3$. Similarly, we can check that (3.12) is positive.

From (iii,1) through (iii,6), we have $e_f(F_p) - \mu\chi_f(F_p) \geq 0$. Thus, Theorem 3.1 (3) follows. $\square$


Example 3.13. There exist singular fiber germs F_p such that $K_f^2(F_p) = (12 - \mu)\chi_f(F_p)$ and we can classify them.

(i) Assume that $n \geq 4$, or $n = 3$ and $g = 4$. Consider the situation that Γ_p is smooth and F_p is obtained the following sequence of singularity diagrams associated with Γ_p (cf. [8]):




Then we can write $F_p = A_0 + \sum_{i=1}^n (A_{1,i} + \dots + A_{l-1,i}) + A_l$, $p_a(A_0) = 0$, $g(A_{k,i}) = 0$ for $k = 1, \dots, l-1$ and $g(A_l) = (r/2 - 1)(n-1)$ (note that A_0 may not be irreducible). This singular fiber satisfies $K_f^2(F_p) = (12 - \mu)\chi_f(F_p)$. Indeed, $\alpha_k(F_p) = 0$ for $k = 0, 1, \dots, r/n - 1$, $\alpha_{r/n}(F_p) = l$, $\varepsilon(F_p) = 0$ and $\chi_\varphi(F_p) = 0$. Thus $\chi_f(F_p) = r(n-1)(n+1)l/12n$, $e_f(F_p) = nl$ and then $e_f(F_p)/\chi_f(F_p) = 12n^2/r(n-1)(n+1) = \mu$. We can see from the proof of Theorem 3.1 that any singular fiber F_p satisfying $K_f^2(F_p) = (12 - \mu)\chi_f(F_p)$ is obtained in this way.

(ii) Assume that $n = 3$ and $g > 4$. Consider the situation that Γ_p is smooth and F_p is obtained the following sequence of singularity diagrams associated with Γ_p :



Then we can write $F_p = A_0 + 2A_1 + A_3$, $p_a(A_0) = 4$, $g(A_1) = 1$ and $g(A_2) = r - 5$. This singular fiber satisfies $K_f^2(F_p) = (12 - \mu)\chi_f(F_p)$. Indeed, $\alpha_0(F_p) = 1$, $\alpha_1(F_p) = l$, $\alpha_{r/3-1}(F_p) = 3$, $\alpha_k(F_p) = 0$ for $k \neq 1, r/3 - 1$, $\varepsilon(F_p) = 2$ and $\chi_\varphi(F_p) = 0$. Thus $\chi_f(F_p) = (4r - 13)/6$, $e_f(F_p) = 4$ and then $e_f(F_p)/\chi_f(F_p) = 24/(4r - 13) = \mu$. We can see from the proof of Theorem 3.1 that any singular fiber F_p satisfying $K_f^2(F_p) = (12 - \mu)\chi_f(F_p)$ is obtained in this way.

(iii) Assume that $n = 2$ and $g \geq 3$. Consider the situation that Γ_p is smooth and F_p is obtained the following sequence of singularity diagrams associated with Γ_p :



Then we can write $F_p = A_0 + A_1 + \cdots + A_l$, $p_a(A_0) = 2$, $g(A_k) = 0$ for $k = 1, \dots, l-1$ and $g(A_l) = r/2 - 1$. This singular fiber satisfies $K_f^2(F_p) = (12 - \mu)\chi_f(F_p)$. Indeed, $\alpha_{r/2-1}(F_p) = \alpha_{r/2}(F_p) = l$, $\alpha_k(F_p) = 0$ for $k \neq r/2 - 1, r/2$, $\varepsilon(F_p) = l$ and $\chi_\varphi(F_p) = 0$. Thus $\chi_f(F_p) = (r-2)l/4$, $e_f(F_p) = l$ and then $e_f(F_p)/\chi_f(F_p) = 4/(r-2) = \mu$. We can see from the proof of Theorem 3.1 that any singular fiber F_p satisfying $K_f^2(F_p) = (12 - \mu)\chi_f(F_p)$ is obtained in this way.

From Theorem 3.1 and Example 3.13, we can characterize primitive cyclic covering fibrations of type $(g, 1, n)$ whose slope attains the upper bound in Theorem 3.1.

Corollary 3.14. *Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(g, 1, n)$. Then the slope λ_f attains the upper bound in Theorem 3.1 if and only if any singular fiber of f is as in Example 3.13.*

4. UPPER BOUND OF THE SLOPE: THE CASE OF TYPE $(g, 0, 3)$, $(g, 0, 2)$

Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(g, 0, n)$. We freely use notations in the previous section. Since $\tilde{\varphi}: \widetilde{W} \rightarrow B$ is a ruled surface, a relatively minimal model of it is not unique. By performing elementary transformations, we can choose a standard one:

Lemma 4.1 (cf. [7], [18]). *There exists a relatively minimal model $\varphi: W \rightarrow B$ of $\tilde{\varphi}$ such that if $n = 2$ and g is even, then*

$$\text{mult}_x(R) \leq \frac{r}{2} = g + 1$$

for all $x \in R$, and otherwise,

$$\text{mult}_x(R_h) \leq \frac{r}{2} = \frac{g}{n-1} + 1$$

for all $x \in R_h$, where R_h denotes the horizontal part of R , that is, the sum of all φ -horizontal components of R .

We take a relatively minimal model $\varphi: W \rightarrow B$ satisfying the inequality in Lemma 4.1. It is easily seen that the value of $\alpha_k(F_p)$ does not depend on the choice of the relatively minimal model of $\tilde{\varphi}$ satisfying Lemma 4.1. Since $\varphi: W \rightarrow B$ is a relatively minimal ruled surface, we have $K_\varphi^2 = 0$ and $R^2 = -rK_\varphi R$. Combining these equalities with (1.8), (1.9), (1.10) and (1.11), we get the following lemma:

Lemma 4.2 (cf. [7]). *The following equalities hold.*

$$\begin{aligned} K_f^2 &= \frac{n-1}{r-1} \left(\frac{(n-1)r-2n}{n} (\alpha_0 - 2\varepsilon) + (n+1) \sum_{k \geq 1} k(-nk+r) \alpha_k \right) - n \sum_{k \geq 1} \alpha_k + \varepsilon. \\ \chi_f &= \frac{n-1}{12(r-1)} \left(\frac{(2n-1)r-3n}{n} (\alpha_0 - 2\varepsilon) + (n+1) \sum_{k \geq 1} k(-nk+r) \alpha_k \right). \\ e_f &= (n-1)\alpha_0 + n \sum_{k \geq 1} \alpha_k - (2n-1)\varepsilon. \end{aligned}$$

For $p \in B$, we put

$$\begin{aligned}
K_f^2(F_p) &= \frac{n-1}{r-1} \left(\frac{(n-1)r-2n}{n} (\alpha_0(F_p) - 2\varepsilon(F_p)) + (n+1) \sum_{k \geq 1} k(-nk+r) \alpha_k(F_p) \right) \\
&\quad - n \sum_{k \geq 1} \alpha_k(F_p) + \varepsilon(F_p), \\
\chi_f(F_p) &= \frac{n-1}{12(r-1)} \left(\frac{(2n-1)r-3n}{n} (\alpha_0(F_p) - 2\varepsilon(F_p)) + (n+1) \sum_{k \geq 1} k(-nk+r) \alpha_k(F_p) \right), \\
e_f(F_p) &= (n-1) \alpha_0(F_p) + n \sum_{k \geq 1} \alpha_k(F_p) - (2n-1) \varepsilon(F_p).
\end{aligned}$$

In [7], we have established the slope equality for a primitive cyclic covering fibration of type $(g, 0, n)$ by using the above representations of K_f^2 , χ_f and e_f , and an upper bound of its slope when $n \geq 4$. In this section, we show that a more careful study as in the previous section gives us an upper bound of the slope when $n = 2, 3$. Namely, we show the following theorem:

Theorem 4.3. (1) *Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(4, 0, 3)$. Then we have*

$$K_f^2 \leq \frac{129}{17} \chi_f$$

with the equality holding if and only if any singular fiber of f is a triple fiber.

(2) *Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(g, 0, 3)$. Assume that $g > 4$, or $g = 4$ and f has no triple fibers. Then we have*

$$K_f^2 \leq \left(12 - \frac{72(r-1)}{4r^2 - 15r + 27 - 36\delta} \right) \chi_f,$$

where $\delta = \begin{cases} 0, & \text{if } r \in 6\mathbb{Z}, \\ 1, & \text{if } r \notin 6\mathbb{Z}. \end{cases}$

(3) *Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(g, 0, 2)$, i.e., a relatively minimal hyperelliptic fibration of genus g . Then we have*

$$K_f^2 \leq \left(12 - \frac{4(2g+1)}{g^2 - 1 + \delta} \right) \chi_f,$$

where $\delta = \begin{cases} 0, & \text{if } g \text{ is odd}, \\ 1, & \text{if } g \text{ is even}. \end{cases}$

Remark 4.4. Theorem 4.3 (3) was shown by Xiao in [19] and the upper bound is known to be sharp [15].

Recall the following lemma from [7] (cf. Lemmas 3.4, 3.5 and 3.6):

Lemma 4.5. *The following hold:*

- (1) $\iota(F_p) = j(F_p) - \eta_p$.
- (2) $\alpha_0^+(F_p) \geq (n-2)(j(F_p) - \eta_p + 2\kappa(F_p)) + \delta_{n=2} \sum_{k \geq 1} 2k\alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co}}(F_p)$.
- (3) $\sum_{k \geq 1} \alpha_k''(F_p) = \sum_{a \geq 1} (an-2)j_{0,a}(F_p) + 2\eta_p - \kappa(F_p)$.

Lemma 4.6. *The following hold.*

- (1) *If $n = 3$, then we have $j_{0,1}(F_p) \leq 2\eta_p + 1 + \sum_{a \geq 2} (2a-1)j_{0,a}(F_p)$, with the equality holding only if F_p is a triple fiber, $r \in 9\mathbb{Z} + 6$, $j(F_p) = j_{0,1}(F_p)$, $\kappa(F_p) = 1$ and*

$$\alpha_0^+(F_p) \geq \frac{5(r-6)}{9} + j(F_p) - \eta_p + 2\kappa(F_p).$$

- (2) *If $n = 2$, then*

$$\sum_{k \geq 1} \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},1}(F_p) \leq \widehat{\eta}_p + 2\eta'_p + \sum_{a \geq 3} (a-2)j_{0,a}(F_p) - j'_{0,1}(F_p).$$

Proof. Suppose that $n = 3$. By the same argument as in the proof of Lemma 3.10, we have

$$j''_{0,1}(F_p) \leq \overline{\eta}_p + \eta''_p + \sum_{a \geq 1} 2aj'_{0,a}(F_p) + \sum_{a \geq 2} (2a-1)j''_{0,a}(F_p).$$

Hence, if $\Gamma_p \not\subset R$, we have

$$j_{0,1}(F_p) \leq 2\eta_p + \sum_{a \geq 2} (2a-1)j_{0,a}(F_p).$$

If $\Gamma_p \subset R$ and Γ_p does not contribute to $j'_{0,1}(F_p)$, we have

$$j_{0,1}(F_p) \leq 2\eta_p - 1 + \sum_{a \geq 2} (2a-1)j_{0,a}(F_p),$$

since $j''_{0,1}(F_p) = j_{0,1}(F_p)$, $\sum_{a \geq 1} 2aj'_{0,a}(F_p) + \sum_{a \geq 2} (2a-1)j''_{0,a}(F_p) = \sum_{a \geq 2} (2a-1)j_{0,a}(F_p) + 1$ and $\overline{\eta}_p \leq \eta''_p \leq \eta_p - 1$. If $\Gamma_p \subset R$ and Γ_p contributes to $j'_{0,1}(F_p)$, we have

$$j_{0,1}(F_p) \leq 2\eta_p + 1 + \sum_{a \geq 2} (2a-1)j_{0,a}(F_p),$$

since $j''_{0,1}(F_p) = j_{0,1}(F_p) - 1$, $\sum_{a \geq 1} 2aj'_{0,a}(F_p) + \sum_{a \geq 2} (2a-1)j''_{0,a}(F_p) = \sum_{a \geq 2} (2a-1)j_{0,a}(F_p) + 2$ and $\overline{\eta}_p \leq \eta''_p \leq \eta_p - 1$. If $j_{0,1}(\overline{F}_p) = 2\eta_p + 1 + \sum_{a \geq 2} (2a-1)j_{0,a}(F_p)$, then $\overline{\eta}_p = \eta_p - 1$ and the sequence of singularity diagrams associated with $D^t(p)$ containing the proper transform of the fiber $\Gamma_p = C^{t,1}$ is the following:

$$\begin{array}{|c|} \hline n_1 \\ \hline m_2 \\ \hline m_1 \\ \hline \end{array} \\ \mathcal{D}^{t,1}$$

$$\begin{array}{|c|} \hline n_2 \\ \hline m_2 \\ \hline \end{array} \\ \mathcal{D}^{t,2}$$

$$\begin{array}{|c|c|} \hline n_1 & n_2 \\ \hline \end{array} \\ \mathcal{D}^{t,3}$$

where $n_i \in 3\mathbb{Z}$, $m_i \in 3\mathbb{Z} + 1$ and $C^{t,k}$ is the exceptional curve obtained by the blow-up at the singularity with multiplicity m_{k-1} for $k = 2, 3$. From Lemma 1.9, we have $r+5 = m_1+m_2+n_1$, $m_1+3 = m_2+n_2$ and $m_2+2 = n_1+n_2$. In particular, we get $r = 3(n_1+n_2-4) \in 9\mathbb{Z}+6$. Since $\bar{\eta}_p = \eta_p''$, the sequence of singularity diagrams associated with another $D^t(p)$ is the following from Example 1.15 (2):

$$\begin{array}{c} \boxed{\begin{array}{c} n_1 \\ m_1 \end{array}} \\ \mathcal{D}^{t,1} \end{array} \qquad \begin{array}{c} \boxed{\begin{array}{c} n_2 \\ n_1 \end{array}} \\ \mathcal{D}^{t,2} \end{array} \quad \text{or} \quad \begin{array}{c} \boxed{\begin{array}{cc} n_1 & n_2 \end{array}} \\ \mathcal{D}^{t,2} \end{array}$$

where $n_i \in 3\mathbb{Z}$, $m_i \in 3\mathbb{Z} + 1$ and $C^{t,2}$ is the exceptional curve obtained by the blow-up at the singularity with multiplicity m_1 . From Lemma 1.9, we have $m+3 = m_1+n_1$ and $m_1+2 = n_1+n_2$, where m is the multiplicity of R at the point to which $C^{t,1}$ is contracted. Clearly, we have $j(F_p) = j_{0,1}(F_p) = 2\eta_p + 1$, $\kappa(F_p) = 1$. Moreover, we see that F_p is a triple fiber since the multiplicity of the fiber over p along the exceptional curve obtained by the blow-up at a singularity of type $3\mathbb{Z}$ is 3-multiple. Hence, we get $\alpha_0^+(F_p) \geq 2r/3$. To prove (1), it is sufficient to show that $2r/3 - (j(F_p) - \eta_p + 2\kappa(F_p)) \geq 5(r-6)/9$, i.e., $\eta_p \leq (r+3)/9$. To prove this, we will show that for any $n_1 \in 3\mathbb{Z}_{>0}$, the number of singularities of type $3\mathbb{Z}+1$ over a singularity with multiplicity n_1 is less than or equal to $2n_1/3 - 2$ by induction on $n_1/3$. If $n_1 = 3$, the claim is obvious. We assume that $n_1 > 3$ and the claim holds true until $n_1 - 3$. Let x be a singularity of R with multiplicity n_1 . Let $x_1, \dots, x_l$ be all singularities of type $3\mathbb{Z} + 1$ over x which is not over any singularity of type $3\mathbb{Z} + 1$ over x . Let m_i denotes the multiplicity of R at x_i for $i = 1 \dots, l$. Then, the sequence of singularity diagrams associated with $D^t(p)$ obtained by the blow-up at x_i is the following:

$$\begin{array}{c} \boxed{\begin{array}{c} n_{i,1} \\ m_{i,1} \end{array}} \\ \mathcal{D}^{t,1} \end{array} \qquad \begin{array}{c} \boxed{\begin{array}{c} n_{i,2} \\ n_{i,1} \end{array}} \\ \mathcal{D}^{t,2} \end{array} \quad \text{or} \quad \begin{array}{c} \boxed{\begin{array}{cc} n_{i,1} & n_{i,2} \end{array}} \\ \mathcal{D}^{t,2} \end{array}$$

where $n_{i,j} \in 3\mathbb{Z}$, $m_{i,j} \in 3\mathbb{Z} + 1$ satisfy that $m_i+3 = m_{i,1}+n_{i,1}$, $m_{i,1}+2 = n_{i,1}+n_{i,2}$ and $C_{t,1}$, $C^{t,2}$ respectively denote the exceptional curves obtained by the blow-up at the singularity x_i , the singularity with multiplicity $m_{i,1}$. By the assumption of induction, the number of singularities of type $3\mathbb{Z} + 1$ over x_i is less than or equal to $2(n_{i,1}+n_{i,2})/3 - 3$. Hence the number of singularities of type $3\mathbb{Z} + 1$ over x is less than or equal to $\sum_{i=1}^l 2(n_{i,1}+n_{i,2})/3 - 2l$. If $l = 1$, then we have $n_1 \geq m_1 + 2 = 2n_{1,1} + n_{1,2} - 3$. Thus, we get

$$\begin{aligned}\frac{2}{3}n_1 - 2 &\geq \frac{2}{3}(2n_{1,1} + n_{1,2}) - 4 \\ &\geq \frac{2}{3}(n_{1,1} + n_{1,2}) - 2.\end{aligned}$$

If $l = 2$, then we have $n_1 \geq m_1 + m_2 + 1 = 2n_{1,1} + n_{1,2} + 2n_{2,1} + n_{2,2} - 9$. Thus, we get

$$\begin{aligned}\frac{2}{3}n_1 - 2 &\geq \frac{2}{3}(2n_{1,1} + n_{1,2} + 2n_{2,1} + n_{2,2}) - 8 \\ &\geq \frac{2}{3}(n_{1,1} + n_{1,2} + n_{2,1} + n_{2,2}) - 4.\end{aligned}$$

If $l \geq 3$, then we have $n_1 \geq \sum_{i=1}^l m_i = \sum_{i=1}^l (2n_{i,1} + n_{i,2}) - 5l$. Thus, we get

$$\begin{aligned}\frac{2}{3}n_1 - 2 &\geq \frac{2}{3} \sum_{i=1}^l (2n_{i,1} + n_{i,2}) - \frac{10}{3}l - 2 \\ &\geq \frac{2}{3} \sum_{i=1}^l (n_{i,1} + n_{i,2}) - 2l\end{aligned}$$

with equality holds only if $l = 3$, $n_1 = m_1 + m_2 + m_3$ and $n_{i,1} = 3$ for any i . Hence, the number of singularities of type $3\mathbb{Z} + 1$ over a singularity with multiplicity n_1 is less than or equal to $2n_1/3 - 2$. Now, $r = 3(n_1 + n_2 - 4)$ and the number of singularities of type $3\mathbb{Z} + 1$ over p is less than or equal to $2(n_1 + n_2)/3 - 2$. Namely, we have

$$j(F_p) - 1 = j''_{0,1}(F_p) \leq \frac{2}{9}r + \frac{2}{3}.$$

Hence, we have $\eta_p = (j(F_p) - 1)/2 = (r + 3)/9$, which is the desired inequality.

Suppose that $n = 2$. By the same argument as in the proof of Lemma 3.10, we have

$$\sum_{k \geq 1} \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},1}(F_p) \leq \widehat{\eta}_p + \sum_{a \geq 1} a j'_{0,a}(F_p) + \sum_{a \geq 3} (a - 2) j''_{0,a}(F_p).$$

Since $\sum_{a \geq 1} a j'_{0,a}(F_p) + \sum_{a \geq 3} (a - 2) j''_{0,a}(F_p) = 2\eta'_p + \sum_{a \geq 3} (a - 2) j_{0,a}(F_p) - j'_{0,1}(F_p)$, the claim (2) follows. $\square$

Lemma 4.7 (cf. Lemma 3.9). *If $n = 2$, then we have*

$$\kappa(F_p) \leq \frac{2}{3} \sum_{a \geq 2} (a - 1) j_{0,a}(F_p).$$

Proof. It is sufficient to show that

$$\kappa^t(F_p) \leq \frac{2}{3} \sum_{a \geq 2} (a - 1) j_{0,a}^t(F_p)$$

for any t . If $\kappa^t(F_p) = 0$, then it is clear. Thus, we may assume $\kappa^t(F_p) > 0$. Then clearly we have $j^t(F_p) \geq \kappa^t(F_p) + 2$. Since any blow-up at a $(t, 2)$ -vertical type singularity contributes -2 to the number $\sum_{k \geq 1} (L^{t,k})^2 = -\sum_{a \geq 1} 2aj_{0,a}^t(F_p)$ and Γ_p contains no 2-vertical type singularity, we get

$$-\sum_{a \geq 1} 2aj_{0,a}^t(F_p) \leq -j^t(F_p) + (-2)(\iota^t(F_p) + \kappa^t(F_p)) = -(3j^t(F_p) + 2\kappa^t(F_p) - 2).$$

Combining this equality with $j^t(F_p) \geq \kappa^t(F_p) + 2$, we have

$$\sum_{a \geq 1} 2aj_{0,a}^t(F_p) \geq 2j^t(F_p) + 3\kappa^t(F_p).$$

Hence we get $\kappa^t(F_p) \leq \frac{2}{3} \sum_{a \geq 2} (a-1)j_{0,a}^t(F_p)$. □

Now, we are ready to prove Theorem 4.3.

Proof of Theorem 4.3. Let $f: S \rightarrow B$ be a primitive cyclic covering fibration of type $(g, 0, n)$. For any $p \geq B$ and $\mu \geq 0$, we have from Lemma 4.5 (3) that

$$\begin{aligned} & e_f(F_p) - \mu\chi_f(F_p) \\ &= (n-1)\alpha_0(F_p) + n \sum_{k \geq 1} \alpha_k(F_p) - (2n-1)\varepsilon(F_p) \\ & \quad - \mu' \left(\frac{r(2n-1) - 3n}{n} \alpha_0(F_p) + (n+1) \sum_{k \geq 1} (-nk^2 + rk) \alpha_k(F_p) - \frac{2(r(2n-1) - 3n)}{n} \varepsilon(F_p) \right) \\ &= A_n \alpha_0(F_p) + \sum_{k \geq 1} (Q(k) + B_n) \alpha_k(F_p) - (2A_n + 1) \varepsilon(F_p) \\ &= A_n \alpha_0^+(F_p) + \sum_{k \geq 1} (Q(k) + B_n) \alpha'_k(F_p) + \sum_{k \geq 1} (Q(k) + B_n) \alpha''_k(F_p) - 2A_n j(F_p) - j_{0,1}(F_p) \\ &= A_n \alpha_0^+(F_p) + \sum_{k \geq 1} (Q(k) + B_n) \alpha'_k(F_p) + \sum_{k \geq 1} Q(k) \alpha''_k(F_p) \\ & \quad + \sum_{a \geq 1} ((an - 2)B_n - 2A_n - \delta_{a=1}) j_{0,a}(F_p) + 2B_n \eta_p - B_n \kappa(F_p), \end{aligned}$$

where

$$\mu' = \frac{n-1}{12(r-1)} \mu, \quad A_n = n-1 - \frac{r(2n-1) - 3n}{n} \mu', \quad B_n = n - \frac{(n+1)(r^2 - \delta n^2)}{4n} \mu'$$

and

$$Q(k) = \mu' \left(n(n+1) \left(k - \frac{r}{2n} \right)^2 - \frac{n(n+1)\delta}{4} \right) \geq 0.$$

Assume that $A_n \geq 0$ and $B_n \geq 0$. From Lemma 4.5 (2), $e_f(F_p) - \mu\chi_f(F_p)$ is greater than or equal to

$$\begin{aligned}
& \left((n-2)(j(F_p) - \eta_p + 2\kappa(F_p)) + \delta_{n=2} \sum_{k \geq 1} 2k\alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co}}(F_p) \right) A_n + \sum_{k \geq 1} Q(k)\alpha_k''(F_p) \\
& + \sum_{a \geq 1} ((an-2)B_n - 2A_n - \delta_{a=1})j_{0,a}(F_p) + 2B_n\eta_p - B_n\kappa(F_p) \\
& = \delta_{n=2} \sum_{k \geq 1} 2k\alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co}}(F_p)A_n + \sum_{a \geq 1} ((n-4)A_n + (an-2)B_n - \delta_{a=1})j_{0,a}(F_p) \\
(4.1) \quad & + \sum_{k \geq 1} Q(k)\alpha_k''(F_p) + (2B_n - (n-2)A_n)\eta_p + (2(n-2)A_n - B_n)\kappa(F_p).
\end{aligned}$$

(i) We first assume that $n = 3$.

(i,1) Assume that $j_{0,1}(F_p) \leq 2\eta_p + \sum_{a \geq 2} (2a-1)j_{0,a}(F_p)$. We put

$$\mu = \frac{72(r-1)}{4r^2 - 15r + 27 - 36\delta}.$$

Then, we have $A_3 > 0$ and $B_3 > 0$. Since the coefficient $-A_3 + 2B_3$ of η_p in (4.1) is also positive, (4.1) is greater than or equal to

$$(4.2) \quad \sum_{a \geq 2} \left(\left(a - \frac{3}{2} \right) A_3 + (a-1)B_3 \right) j_{0,a}(F_p) + \left(-\frac{3}{2}A_3 + 2B_3 - 1 \right) j_{0,1}(F_p) + (2A_3 - B_3)\kappa(F_p).$$

By the definition of μ , we have $-\frac{3}{2}A_3 + 2B_3 - 1 = 0$ and $2A_3 - B_3 > 0$. Thus (4.2) is non-negative.

(i,2) Assume that $j_{0,1}(F_p) = 2\eta_p + 1 + \sum_{a \geq 2} (2a-1)j_{0,a}(F_p)$ and $r > 6$. We also put

$$\mu = \frac{72(r-1)}{4r^2 - 15r + 27 - 36\delta}.$$

From Lemma 4.6 (1), we have

$$\begin{aligned}
& e_f(F_p) - \mu\chi_f(F_p) \\
& \geq \frac{5(r-6)}{9}A_3 + (-A_3 + B_3 - 1)j_{0,1}(F_p) + (-A_3 + 2B_3)\eta_p + (2A_3 - B_3) \\
& = \frac{5(r-6)}{9}A_3 + \left(-\frac{3}{2}A_3 + 2B_3 - 1\right)j_{0,1}(F_p) + \frac{5}{2}A_3 - 2B_3 \\
& \geq \frac{15}{2}A_3 - 2B_3 \\
& > 0.
\end{aligned}$$

(i,3) Assume that $j_{0,1}(F_p) = 2\eta_p + 1 + \sum_{a \geq 2} (2a - 1)j_{0,a}(F_p)$ and $r = 6$. We put $\mu = \frac{129}{17}$. From Lemma 4.6 (1) and $j_{0,1}(F_p) = 3$, we have

$$e_f(F_p) - \mu\chi_f(F_p) \geq -2A_3 + 4B_3 - 3 = 0,$$

where the last equality follows by the definition of μ .

(ii) We assume that $n = 2$. Put

$$\mu = \frac{4(2g + 1)}{g^2 - 1 + \delta}.$$

Then we have $2B_2 - 2A_2 - 1 - 6(1 - \delta)\mu' = 0$, $A_2 > 0$ and $B_2 \geq 1$. From (1.12) and (1.14), (4.1) is equal to

$$\begin{aligned} & \sum_{k \geq 1} Q(k)\alpha_k''(F_p) + \sum_{a \geq 2} (-2A_2 + (2a - 2)B_2)j_{0,a}(F_p) - B_2\kappa(F_p) \\ & + (2B_2 - 2A_2 - 1) \sum_{k \geq 1} \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{tr}}(F_p) + \sum_{k \geq 1} (2(k - 1)A_2 + 2B_2 - 1) \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},0}(F_p) \\ & + \sum_{k \geq 1} (2(k - 1)A_2 - 1) \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},1}(F_p) - (2A_2 + 1)j_{0,1}'(F_p) + 2B_2(\eta_p' + \widehat{\eta}_p). \end{aligned}$$

Using Lemma 4.6 (2) for $-\sum_{k \geq 1} \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},1}(F_p)$, (4.1) is greater than or equal to

$$\begin{aligned} & \sum_{k \geq 1} Q(k)\alpha_k''(F_p) + \sum_{a \geq 2} (-2A_2 + (2a - 2)B_2 - (a - 2))j_{0,a}(F_p) - B_2\kappa(F_p) \\ & + (2B_2 - 2A_2 - 1) \sum_{k \geq 1} \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{tr}}(F_p) + \sum_{k \geq 1} (2(k - 1)A_2 + 2B_2 - 1) \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},0}(F_p) \\ (4.3) \quad & + \sum_{k \geq 1} 2(k - 1)A_2 \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},1}(F_p) - 2A_2 j_{0,1}'(F_p) + (2B_2 - 2)\eta_p' + (2B_2 - 1)\widehat{\eta}_p. \end{aligned}$$

By the definition of μ , we see that

$$(4.4) \quad \sum_{k \geq 1} Q(k)\alpha_k''(F_p) + (2B_2 - 2A_2 - 1) \sum_{k \geq 1} \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{tr}}(F_p) \geq 0$$

with the equality holding if and only if $\alpha_{(g-1)/2}''(F_p) = \alpha_{(g \rightarrow g)}^{\text{tr}}(F_p)$ and $\alpha_k''(F_p) = 0$ for $k \neq (g - 1)/2$ when g is odd, or $\alpha_{g/2}''(F_p) = \alpha_{(g+1 \rightarrow g+1)}^{\text{tr}}(F_p)$ and $\alpha_k''(F_p) = 0$ for $k \neq g/2$ when g is even. If $j_{0,1}'(F_p) = 1$, then g is odd and $\Gamma_p \subset R$ contains a singularity of type $(g + 2 \rightarrow g + 2)$ since Γ_p is blown up just twice and Lemma 4.1. Thus, Γ_p contributes to $\alpha_{(g+2 \rightarrow g+2)}^{\text{co},1}(F_p)$ and then we have

$$(4.5) \quad \sum_{k \geq 1} 2(k - 1)A_2 \alpha_{(2k+1 \rightarrow 2k+1)}^{\text{co},1}(F_p) - 2A_2 j_{0,1}'(F_p) \geq 0.$$

From Lemma 4.7 and

$$-2A_2 + \frac{4}{3}(a - 1)B_2 - (a - 2) > 0,$$

we have

$$(4.6) \quad \sum_{a \geq 2} (-2A_2 + (2a - 2)B_2 - (a - 2)) j_{0,a}(F_p) - B_2 \kappa(F_p) \geq 0.$$

From (4.4), (4.5), (4.6) and $B_2 \geq 1$, (4.3) is non-negative. $\square$

REFERENCES

- [1] T. Ashikaga and K. Konno, Global and local properties of pencils of algebraic curves, Algebraic Geometry 2000 Azumino, S. Usui et al. eds, 1-49, Adv. Stud. Pure Math. **36**, Math. Soc. Japan, Tokyo, 2002.
- [2] M.A. Barja, On the slope of bielliptic fibrations, Proc. Amer. Math. Soc. **129** (2001), 1899–1906.
- [3] M. A. Barja and J. C. Naranjo, Extension of maps defined on many fibres, Dedicated to the memory of Fernando Serrano, Collect. Math. **49** (1998), 227–238.
- [4] W. Barth, K. Hulek, C. Peters and A. Van de Ven, *Compact Complex Surfaces*, 2nd Edition, Springer-Verlag, Berlin, 2004.
- [5] Z. Chen and S-L. Tan, Upper bounds on the slope of a genus 3 fibration, Recent progress on some problems in several complex variables and partial differential equations, 65–87., Contemp Math. **400**, Amer. Math. Soc., Providence, RI, 2006.
- [6] M. Cornalba and L. Stoppino, A sharp bound for the slope of double cover fibrations, Michigan Math. J. **56** (2008), 551–561.
- [7] M. Enokizono, Slopes of fibered surfaces with a finite cyclic automorphism, to appear in Michigan Math. J, arXiv:1504.04106[math.AG].
- [8] M. Enokizono, Fibers of cyclic covering fibrations of a ruled surface, arXiv:1505.03904[math.AG].
- [9] P. A. Griffiths and J. Harris, *Principles of Algebraic Geometry*, Wiley Classics Library. John Wiley & Sons Inc., New York (1994).
- [10] A. Kas, On deformations of a certain type of irregular algebraic surface, Amer. J. Math. **90** (1968), 789–804.
- [11] K. Kodaira, A certain type of irregular algebraic surfaces, J. Anal. Math. **19** (1967), 207–215.
- [12] K. Kodaira, On compact complex analytic surfaces II, Ann. of Math. **77** (1963), 563–626.
- [13] K. Konno, Non-hyperelliptic fibrations of small genus and certain irregular canonical surfaces, Ann. Sc. Norm. Sup. Pisa ser. IV, **20** (1993), 575–595.
- [14] K. Konno, Clifford index and the slope of fibered surfaces, J. Alg. Geom. **8** (1999), 207–220.
- [15] X.-L. Liu and S.-L. Tan, Families of hyperelliptic curves with maximal slopes, Sci. China Math. **56** (2013), 1743–1750.
- [16] S. Matsusaka, Some numerical invariants of hyperelliptic fibrations, J. Math. Kyoto Univ. **30** (1990), 33–57.
- [17] G. Xiao, Fibered algebraic surfaces with low slope, Math. Ann. **276** (1987), 449–466.
- [18] G. Xiao, π_1 of elliptic and hyperelliptic surfaces, Internat. J. Math. **2** (1991), 599–615.
- [19] G. Xiao, *Fibrations of Algebraic Surfaces* (in Chinese), Shanghai Publishing House of Science and Technology, 1992.

MAKOTO ENOKIZONO, DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN

E-mail address: m-enokizono@cr.math.sci.osaka-u.ac.jp